

Optimal Routing in Output-Queued Flexible Server Systems

Alexander L. Stolyar
Bell Labs, Lucent Technologies
Murray Hill, NJ 07974
`stolyar@research.bell-labs.com`

May 3, 2004

Abstract

We consider a queueing system with multi-type customers and non-homogeneous flexible servers, in the heavy traffic asymptotic regime and under a *complete resource pooling* (CRP) condition. For the *input-queued* (IQ) version of such a system (with customers being queued at the system “entrance,” one queue per each type), it was shown in [12] that a simple parsimonious $G\mu$ scheduling rule is optimal in that it asymptotically minimizes the system *customer workload* and some strictly convex queueing costs. In this paper we consider a different - *output-queued* (OQ) - version of the model, where each arriving customer must be assigned to one of the servers immediately upon arrival. (This constraint can be interpreted as immediate *routing* of each customer to one of the “output queues,” one queue per each server.) Consequently, the space of controls allowed for an OQ-system is a subset of that for the corresponding IQ-system.

We introduce the *MinDrift* routing rule for OQ-systems (which is as simple and parsimonious as $G\mu$), and show that this rule, in conjunction with arbitrary work-conserving disciplines at the servers, has asymptotic optimality properties analogous to those $G\mu$ -rule has for IQ-systems. A key element of the analysis is the notion of system *server workload*, which, in particular, majorizes customer workload. We show that (i) MinDrift-rule asymptotically minimizes server workload process among all OQ-system disciplines and (ii) this minimal process matches the minimal possible customer workload process in the corresponding IQ-system. As a corollary, MinDrift asymptotically minimizes customer workload among all disciplines in either the OQ- or IQ-system.

Key words and phrases: Flexible servers, MinDrift, optimal dynamic routing, load balancing, queueing networks, heavy traffic (diffusion) limit, resource pooling

1 Introduction

1.1 The problem.

We consider a queueing system with multiple customer flows (types) $i = 1, \dots, I$ and non-homogeneous flexible servers $j = 1, \dots, J$. This means that the mean service time μ_{ij}^{-1} of a type i customer by server j depends on both the customer type and the server. We study the “heavy traffic” asymptotic regime, when the system is close to be “critically loaded,” and assume that a certain *Complete Resource Pooling* (CRP) condition holds. Associated with the CRP condition, is the notion of system *workload*, which in this paper is called system *customer workload*.

The “input-queued” (IQ) version of the model (see [8, 3, 9, 20, 12]) is such that arriving customers are placed in “input” queues, one queue per each type i , where they await for service without being pre-assigned to any particular server until they are actually “taken for service” by one of them. It is shown in [12], that an IQ-system can be asymptotically optimally controlled in heavy traffic (and under CRP condition) by a very simple and parsimonious $Gc\mu$ scheduling rule, which, in particular, minimizes customer workload among virtually all service disciplines.

In this paper we consider a different - “output-queued” (OQ) - version of the model, where each arriving customer must be assigned (or *routed*) to one of the servers immediately upon arrival. (This can be viewed as immediate routing of each arriving customer to one of the “output” queues, one per each server.) Such models arise in various applications, including wireless networks, manufacturing systems, call centers. A wireless application example is a system where data packets (“customers”) need to be delivered to multiple mobile users/destinations (which determine “customer types”) via a set of transmitters (“servers”); and transmission (“service”) rates depend on the (different) channel qualities between different transmitters and users.

Due to the above *immediate routing* (IR) constraint, the space of controls allowed for an OQ-system is a (strict) subset of that for the corresponding IQ-system. ($Gc\mu$ is not a valid discipline for OQ-systems.) A natural question is: “Is it possible to control OQ-system in heavy traffic as efficiently as the corresponding IQ-system could be controlled? For example, are there OQ-system controls, which are as parsimonious as $Gc\mu$, but still able to minimize the system customer workload?” The results of this paper demonstrate that the answer to the latter question is ‘yes’ - we introduce the *MinDrift* routing rule, which, in particular, minimizes the system customer workload in heavy traffic. We will describe our results shortly, after a brief literature review.

1.2 Previous work.

The IQ-version of our system (with non-homogeneous flexible servers) and the definition of the heavy traffic asymptotic regime for it (in terms of a certain linear program) were introduced in [8] (for a special two-server system) and in [9, 20] (for the general setting). For the system in heavy traffic, these papers also define the CRP condition and associated with it notion of *customer workload*, defined as $X(t) = \sum_i \nu_i^* Q_i(t)$, $t \geq 0$, where $Q_i(t)$ is the type i queue length at time t , and $\nu_i^* > 0$ is the fixed constant called *workload contribution* of a type i customer. Papers [8, 9] propose *discrete-review* scheduling policies which, in the heavy traffic and under CRP condition, asymptotically minimize customer workload and a linear holding costs. In papers [3, 20], *continuous-review threshold* policies are proposed which are asymptotically optimal, in the same sense and also under CRP condition. (The asymptotic optimality proofs are given for a special two-server system.) The common feature of discrete-review and continuous-review threshold policies is that they require a priori knowledge of the flows' mean arrival rates λ_i .

In [12], it is proved that a very simple *generalized cμ* ($Gc\mu$) scheduling rule (which, in particular, does *not* require the knowledge of arrival rates λ_i) asymptotically minimizes customer workload and strictly convex holding costs in a general IQ-system under CRP condition. Moreover, in the limit, the (appropriately rescaled) queue length vector-process $(Q_1(t), \dots, Q_I(t))$ exhibits *state space collapse* (SSC) - it "lives" on a one-dimensional manifold. (The results of [12] are closely related to earlier results in [14] for a discrete time *generalized switch* model. Also, they generalize the earlier $Gc\mu$ optimality results for a single-server system.) Following [14], paper [12] provides equivalent (geometric) characterization of the CRP condition. Namely, let \mathcal{M} be the system *service rate region*, which is roughly the set of all vectors, representing feasible long-term average service rates the system is capable of *jointly* providing to different types. Then, the vector $\nu^* = (\nu_1^*, \dots, \nu_I^*)$ of workload contributions is the *unique* (up to scaling) outer normal vector to the boundary of \mathcal{M} at the point $\lambda = (\lambda_1, \dots, \lambda_I)$.

Most of the previous work on OQ-systems is concentrated on load balancing schemes for systems with homogeneous servers. Much less work has been done on heavy traffic regime in systems with non-homogeneous flexible servers. Probably the first was [10], where a two-server system is considered, resource pooling in heavy traffic is discussed, and threshold-based policies are proposed. In a recent paper [16] a two-server system (different from that in [10]) with exponential service times is considered, and asymptotic optimality of a threshold routing policy is proved, under linear holding costs. We refer reader to [16] for a more extensive review of the previous work on OQ-systems.

1.3 Our results.

In this paper we consider a general OQ-system in heavy traffic regime and under CRP condition. First, we give further equivalent characterization of the CRP condition, which is natural and convenient for the analysis of OQ-systems. Namely, we consider the *server utilization*

region \mathcal{K} , which is the set of potential server utilization vectors, which can be imposed by the input flows with mean rates λ_i or greater. Then, CRP condition, in particular, implies the uniqueness (up to scaling) of the vector normal to \mathcal{K} at the boundary point $(1, \dots, 1)$. The components $\alpha_j^* > 0$ of the vector $\alpha^* = (\alpha_1^*, \dots, \alpha_J^*)$, opposite of the mentioned above normal vector, we call *server workload contributions* of different servers, and we call by system *server workload* the quantity $^u X(t) = \sum_j \alpha_j^* U_j(t)$, where $U_j(t)$ is the *unfinished work* of server j at time t . We establish the relations between customer and server workload contributions, which show that the asymptotic relation $X(t) \leq {}^u X(t)$ between customer workload and server workload exists.

We assume that a strictly convex increasing function $C_j(\cdot)$ is defined for each server j , which is interpreted as the cost rate incurred by the unfinished work on server j .

We introduce two versions of the *MinDrift* routing rule. The *MinDrift(U)* assigns an arriving type i customer to a server

$$j \in \arg \min_j \mu_{ij}^{-1} C'_j(U_j(t)) . \quad (1)$$

(This version may not be practical in many cases, because it assumes exact knowledge of the unfinished work values $U_j(t)$.) The *MinDrift(Q)* rule is the same as *MinDrift(U)*, except $U_j(t)$ in (1) is replaced by the *Q-estimated unfinished work* ${}^q U_j(t) = \sum_m \mu_{mj}^{-1} Q_{mj}(t)$ of server j , where $Q_{ij}(t)$ denotes the number of type i customers in server j queue. (*MinDrift(Q)* is a more practical version.)

Our main result (see Theorems 1 and 2) is that, in OQ-system in the heavy traffic asymptotic limit and under CRP condition, the *MinDrift* rule (either version), in conjunction with virtually arbitrary work-conserving scheduling disciplines at the servers, minimizes (among all service disciplines) the server workload and the instantaneous and cumulative costs corresponding to the cost rate $\sum_j C_j(U_j(t))$. Moreover, in the limit, the (rescaled) unfinished work vector-process $(U_1(t), \dots, U_J(t))$ exhibits SSC such that the vector $(C'_1(U_1(t)), \dots, C'_J(U_J(t)))$ is always proportional to α^* . (This behavior is analogous to that exhibited by IQ-systems in [12, 14], but with the server unfinished works replacing “input” queue lengths, and server workload contributions replacing customer workload contributions.) In addition, the minimal server workload process (attained under *MinDrift*) matches the minimal possible customer workload process in the corresponding IQ-system. As a corollary, *MinDrift(Q)* (asymptotically) minimizes customer workload among all disciplines in either the OQ- or IQ-system (see Theorem 3). In this sense, *MinDrift* rule controls an OQ-system as efficiently as the corresponding IQ-system (allowing a wider class of disciplines) can possibly be controlled.

Essentially as another corollary of the main results, we obtain a necessary condition (Theorem 4) for any OQ-system service discipline to have (asymptotic) workload minimization property. Using this condition we demonstrate (in Section 13) that even some very natural service disciplines in OQ-systems, known to guarantee stability of the queues (if such is feasible at all), do *not* minimize system workload in heavy traffic.

Another contribution of the paper is that, in addition to the CRP condition, we in fact identify and characterize a weaker First-Order-CRP (FO-CRP) condition. The purpose of

doing this is two-fold. First, FO-CRP is sufficient to establish convergence properties of *fluid sample paths*, which arise in the *fluid limit* asymptotic regime and (in addition to being an important step in proving the main heavy traffic results) are of independent interest. Secondly, this clarifies the role of the additional assumption, which strengthens FO-CRP to CRP, in proving the heavy traffic results.

Finally, we would like to point out that, despite the fact that the technical development in this paper is in many ways analogous to that in [14, 12], some parts of it are quite different. In particular, the representation of the server workload process in Sections 9 and 10 (roughly, as a sum of the “driving” and “regulation” processes) is substantially different from the representation of customer workload processes in [14, 12].

1.4 Paper outline.

In Section 2 we set basic notations and conventions. The OQ-system model is formally introduced in Section 3. In Section 4 we define the (two versions of) MinDrift routing rule, discuss its basic intuition and some examples. The definition and characterization of FO-CRP and CRP conditions in terms of an IQ-system are presented in Section 5. The following Section 6 gives an equivalent characterization of FO-CRP and CRP conditions in terms of an OQ-system, and establishes relations between customer and server workload contributions. The heavy traffic asymptotic regime is defined in Section 7, which also contains the definitions of and the relations between customer workload and server workload. Section 8 contains formulations of our main results (Theorems 1-4), described earlier. The analysis of fluid sample paths is the subject of Section 9. Section 10 contain the proof of Theorem 1, regarding the asymptotic optimality of MinDrift(U) rule. The proof of Theorem 2 (regarding the MinDrift(Q) rule) can essentially be reduced to that of Theorem 1 - a detailed outline of this reduction is given in Section 11. Theorem 4 - a necessary condition for asymptotic workload minimization - is proved in Section 12. We conclude in Section 13 with a discussion of the relation between stability and heavy traffic optimality properties of service disciplines.

2 Basic Notation and Conventions

We use the standard notations R and R_+ for the sets of real and real non-negative numbers, respectively; and the not quite standard R_{++} for the set of *strictly* positive real numbers. Corresponding N -times product spaces are denoted R^N , R_+^N , and R_{++}^N . The space R^N is viewed as a standard vector-space, with elements $x \in R^N$ being row-vectors $x = (x_1, \dots, x_N)$. We write simply 0 for the zero vector in R^N , and $\mathbf{1} \doteq (1, 1, \dots, 1)$ for a vector with all unit coordinates. (The dimensions of vectors 0 and $\mathbf{1}$ are either specified explicitly or are clear from the context.)

The scalar product (dot-product) of $x, y \in R^N$, is

$$x \cdot y \doteq \sum_{i=1}^N x_i y_i ;$$

and the norm of x is

$$\|x\| \doteq \sqrt{x \cdot x} .$$

Vector inequalities are to be understood component-wise. As an example, for $\gamma, x \in R^N$, $\gamma < x$ means $\gamma_i < x_i$, $i = 1, \dots, N$. Also

$$\gamma \times x \doteq (\gamma_1 x_1, \dots, \gamma_N x_N) ,$$

and if $\gamma \in R_{++}^N$, we slightly abuse notation by writing

$$1/\gamma \doteq (1/\gamma_1, \dots, 1/\gamma_N) .$$

We denote the minimum and maximum of two real numbers ξ_1 and ξ_2 by $\xi_1 \wedge \xi_2$ and $\xi_1 \vee \xi_2$, respectively.

Let $D([0, \infty), R)$ be the standard Skorohod space of right-continuous left-limit (RCLL) functions, defined on $[0, \infty)$ and taking real values. (See, for example, [7] for the definition of this space and its associated topology and σ -algebra.)

The symbol \xrightarrow{w} denotes convergence in distribution of *random processes* (or other random elements), i.e. weak convergence of their *distributions*. Typically, we consider convergence of processes in $D([0, \infty), R)$, or its N -times product space $D^N([0, \infty), R)$ equipped with product topology and σ -algebra.

The symbol $\xrightarrow{u.o.c.}$ (or the abbreviation *u.o.c.* after a convergence statement) stands for convergence that is *uniform on compact sets*, for *elements* of $D([0, \infty), R)$ or its N -times product $D^N([0, \infty), R)$. For functions with a bounded domain $A \subset R$, the *u.o.c.* convergence means uniform convergence.

We reserve the symbol \Rightarrow for weak convergence of *elements* in the space $D([0, \infty), \bar{R})$; the latter is the space of RCLL functions taking values in the set \bar{R} of real numbers, extended to include the two “infinite numbers” $+\infty$ and $-\infty$ (with the natural topology on \bar{R}). If $h, g \in D([0, \infty), \bar{R})$, then $h \Rightarrow g$ means $h(t) \rightarrow g(t)$ for every $t > 0$ where g is continuous. (Convergence at $t = 0$ is not required.) We shall not need any characterization of the topology on $D([0, \infty), \bar{R})$, beyond the definition of convergence given above.

3 The Model

We consider a queueing system with a finite number I of customer *types*, and a finite number J of flexible *servers*. For notational convenience we use the symbol I also for the set of types $\{1, \dots, I\}$. Similarly, J also denotes the set of servers $\{1, \dots, J\}$.

The arrival process for each type $i \in I$ is a renewal process with the time (from the initial time 0) until the first arrival being $u_i(0)$, and the rest of the interarrival times being an i.i.d. sequence $u_i(n), n = 1, 2, \dots$. Let $\lambda_i = 1/E[u_i(1)] > 0$ denote the arrival rate for type i and $\alpha_i^2 = \text{Var}[u_i(1)]$.

The service times of type i customers by server $j \in J$ form an i.i.d. sequence $v_{ij}(n), n = 1, 2, \dots$. Let $\mu_{ij} = 1/E[v_{ij}(1)] < \infty$ and $\beta_i^2 = \text{Var}[v_{ij}(1)]$. The convention $\mu_{ij} = 0$ is used when server j can not serve type i .

All arrival and service processes are assumed mutually independent.

A version of such flexible (parallel) server model, which received most attention in the previous work (see [8, 9, 20, 3, 12] and references therein) is the *input-queued* (IQ) model. In input-queued model, customers of each type i that await service are waiting in a separate “input” queue i of infinite capacity. This in particular means that customers do not have to be assigned to the servers while waiting in the input queue - such server assignment is (irreversibly) done when the customer is “pulled” for service by one of the servers.

In this paper we concentrate on a different - *output-queued* (OQ) - model, satisfying the following (additional) *immediate routing* (IR) condition:

(IR) *Each new customer arriving in the system must be assigned to one of the servers j immediately upon arrival, and after that the customer can only be served by the server it is assigned to.*

A natural way to interpret (IR)-condition (and this interpretation is in fact the main motivation for the OQ-model) is that, upon arrival, each new customer must be *routed* to one of the servers or, more precisely, into the “output” queue associated with (or, “located at”) that server.

Remark 1. It should be clear that the (IR)-condition defines the *only* difference between an OQ-system and the corresponding IQ-system. Therefore, in general, the class of controls (or, service disciplines) for an OQ-system is a *strict subset* of that for the corresponding IQ-system. For example, the $G\mu$ discipline for IQ-systems, studied in [12], does *not* satisfy (IR)-condition and, consequently, is *not* a valid discipline for OQ-systems.

A *service discipline* in an output-queued system consists of two parts: *routing (server assignment) algorithm* and *scheduling rule* employed by each server (and, generally speaking, depending on the server) to determine which customer to serve from its queue, i.e., among the customers assigned to it.

We will consider the class of service disciplines satisfying (in addition to (IR)) the following condition on the routing algorithm:

(d0) *The realizations of a customer service requirements are not known at the time when routing decision (server assignment) for this customer is made. (The distributions of the service requirements at different servers are known.)*

Sometimes, but not always, we will further restrict the class of service disciplines by imposing the following conditions on the server scheduling rules:

(d1) *Scheduling rule of each server is non-preemptive within each customer type. Namely, a server can not take for service a new customer of type i if it already has another type i customer “in service” (with both elapsed and residual service times being non-zero). Consequently, at any given time, a server can not have in service more than one customer of any given type.*

(d2) *A server does not “know” the realization of a customer service time before the customer service starts.*

Note that conditions (d1) and (d2) do *allow* a server idling (even if it has customers in service) or preemption of service of one customer by another customer but of a *different type*. They also allow server-sharing by several customers but, again, each of a different type.

Remark 2. Note that the class of IQ-system service disciplines, satisfying conditions (d0)-(d2) (but not (IR)), is well defined, and it, first, belongs to the class of disciplines studied in [12], and, second, contains the $G\mu$ discipline (see [12]). Moreover, this class is obviously a superset of the defined above class of OQ-system disciplines satisfying (d0)-(d2) and (IR).

4 The MinDrift Routing Rule

4.1 Notation.

Let $U_j(t)$ denote the *(unfinished) work* of server j at time t , namely, the total amount of unfinished processing time of all customers of all types present in server j queue at time t . We denote by $Q_{ij}(t)$ the number of type i customers in queue j at time t , including those customers whose service already started but not yet completed. The quantity

$${}^qU_j \doteq \sum_i (1/\mu_{ij})Q_{ij}(t)$$

we will call *Q -estimated (unfinished) work* of server j . Finally, by $Q_i(t)$ we will denote the total number of type i customers in the system at time t . In the OQ-system, we always have

$$Q_i(t) = \sum_j Q_{ij}(t) ,$$

but we note that $Q_i(t)$ is well defined for both the OQ- and IQ-systems.

4.2 MinDrift Rule Definition.

Suppose that, for each server j , a cost function $C_j(\zeta)$, $\zeta \geq 0$, is given. Assume that the cost functions have the following properties:

$C_j(\cdot)$ is continuous strictly-increasing convex, with $C_j(0) = 0$;
 moreover, the first derivative $C'_j(\cdot)$ is continuous strictly-increasing, with $C'_j(0) = 0$;
 finally, the second derivative $C''_j(\cdot)$ is strictly positive continuous in the open interval $(0, \infty)$,
 with $C''_j(0) = \lim_{\zeta \downarrow 0} C''_j(\zeta) \geq 0$, where $C''_j(0)$ is either finite or is $+\infty$.

The MinDrift rule routes (assigns) customers to the servers as follows. When a new customer of type i arrives in the system, it is routed to a server j such that

$$j \in \arg \min_{j \in J} C'_j(U_j(t))/\mu_{ij} . \quad (2)$$

Ties are broken arbitrarily: for example, in favor of the smallest index j . Also, by convention, a type i customer can never be routed to a server j if $\mu_{ij} = 0$. (Throughout this paper we also adopt a related convention that any expression involving division by μ_{ij} holds under the additional assumption that $\mu_{ij} > 0$, even if we do not state this explicitly.)

Defined above by (2) is the basic version of MinDrift rule - we will refer to it as a MinDrift(U) rule.

A version of MinDrift rule, with U_j in (2) replaced by ${}^q U_j$, will be called MinDrift(Q) rule. Namely, *MinDrift(Q) rule routes an arriving type i customer to a server j such that*

$$j \in \arg \min_{j \in J} C'_j({}^q U_j(t))/\mu_{ij} . \quad (3)$$

4.3 Nature of MinDrift Rule.

The nature of MinDrift rule is simple - it “myopically” (or, “greedily”) tries to minimize the average drift of the aggregate cost function $\sum_j C_j(U_j(t))$. Indeed, $C'_j(U_j(t))/\mu_{ij}$ (see (2)) approximates the expected increment of the aggregate cost function, caused by routing one type i customer (arrived at time t) to server j ; therefore, by (2), MinDrift(U) routes new arrivals in the way such that the (approximate) expected increment of $\sum_j C_j(U_j(t))$ is minimized. In other words, MinDrift(U) routing tries to *minimize the average rate of increase of $\sum_j C_j(U_j(t))$, due to placement of new work (or, load) to the servers*. Note that, in the OQ-system, the “best” a service discipline can do to maximize the rate at which $\sum_j C_j(U_j(t))$ is decremented due to processing of the unfinished work, is to never idle servers when they have work to do. Thus, MinDrift(U) routing rule (in conjunction with arbitrary work-conserving scheduling rules at the servers) strives to minimize the average drift of the aggregate cost.

The MinDrift(Q) rule is based on the same principle as MinDrift(U), except that instead of using the exact values U_j of unfinished work (which may not be available in many application), it uses their (estimated) average values ${}^q U_j$ (which may be more readily available). As we will demonstrate, in the heavy traffic asymptotic regime we consider, the two versions MinDrift(U) and MinDrift(Q) of the rule are in a certain sense “indistinguishable,” under non-restrictive additional conditions.

The $\text{Gc}\mu$ scheduling rule for IQ-systems, studied in [12], is the rule which myopically tries to minimize the drift of the aggregate cost function $\sum_i C_i(Q_i(t))$ of the queue lengths Q_i , with $C_i(\cdot)$ being cost functions having the same properties as functions $C_j(\cdot)$ defined above. Consequently, $\text{Gc}\mu$ is such that a server j always tries to serve a queue

$$i \in \arg \max_{i \in I} C'_i(Q_i(t)) \mu_{ij},$$

thus *maximizing the average rate at which $\sum_i C_i(Q_i(t))$ is decreased due to departures of served customers*. The $\text{Gc}\mu$ does not - and *cannot* - exercise any control over the rate of increase of $\sum_i C_i(Q_i(t))$ due to new arrivals. Therefore, although both $\text{Gc}\mu$ (in an IQ-system) and MinDrift (in an OQ-system) strive to minimize drifts of certain cost functions, they differ in that they control different system state variables: $\text{Gc}\mu$ controls the rates at which queue lengths Q_i are depleted due to service, and MinDrift controls the rate at which unfinished works U_j are increased due to new arrivals.

4.4 Examples.

Consider a special case when the cost functions have the form $C_j(\zeta) = \gamma_j \zeta^{\eta+1}$, with some fixed $\eta > 0$ and $\gamma_j > 0$. Then MinDrift(U) rule becomes: *route an arriving type i customer to a server*

$$j \in \arg \min_{j \in J} \gamma_j (\eta + 1) \frac{[U_j(t)]^\eta}{\mu_{ij}} ; \quad (4)$$

and similarly for MinDrift(Q).

Consider a special case of the system such that, for any given pair (ij) , we have either $\mu_{ij} = 0$ or $\mu_{ij} = \mu_j > 0$. In other words, the system is such that each server j has a (depending on j) subset of types i which it can serve, but the average service rates of all types within this subset are same and equal to μ_j . (And server cannot serve at all any types i outside the subset.) For this system, the MinDrift(Q) version of (4), with $\eta = 1$, becomes:

$$j \in \arg \min_{j \in J} 2\gamma_j \frac{\sum_i Q_{ij}(t)}{\mu_j^2} . \quad (5)$$

Since parameters $\gamma_j > 0$ can be set arbitrarily, we see from (5) that, *for this special system*, such “popular” routing rules as “Join a server j with the shortest queue”:

$$j \in \arg \min_{j \in J} \sum_i Q_{ij}(t) , \quad (6)$$

and “Join a server j with the smallest expected unfinished work”:

$$j \in \arg \min_{j \in J} \frac{\sum_i Q_{ij}(t)}{\mu_j} , \quad (7)$$

are special cases of MinDrift(Q). (We remind that, in both cases, routing customers to servers where they cannot be served at all is prohibited. Also note that, if we further assume that each server employs first-in-first-out (FIFO) scheduling, then rule (7) is equivalent to the “Join the shortest expected delay” routing rule.)

It should be clear that in a general system, where service rates μ_{ij} depend on both i and j more generally than in the special system described above, the Join-shortest-queue and Join-smallest-expected-unfinished-work routing rules are *not* special cases of MinDrift. Consequently, the heavy traffic optimality properties (which we prove in this paper for MinDrift) may not (and typically *do not*) hold for these rules.

5 Complete Resource Pooling Condition

In this section we present the definition of the Complete Resource Pooling (CRP) condition and related notions and results, which are “oriented towards” the analysis of the IQ-model, and basically follow those in [12]. However, the development in this section is more general than that in Section 5 of [12]. In particular, we consider the notion of First-Order-CRP (which is a weaker form of CRP), and prove some additional properties related to this notion. (The results of this section provide a “reference point” for the next section, which gives an equivalent characterization of First-Order-CRP and CRP conditions “in terms of” the OQ-model.)

Consider an $I \times J$ matrix $\phi = \{\phi_{ij}, i \in I, j \in J\}$, with all $\phi_{ij} \geq 0$. Each element ϕ_{ij} can be interpreted as the average rate at which server j time is allocated to the service of type i customers, in the long run. We do not call elements ϕ_{ij} “fractions” of server j time, because, for the reasons which will become clear in Section 6, it will be convenient for us *not* to assume a priori that $\sum_i \phi_{ij} \leq 1$, or even that $\phi_{ij} \leq 1$.

With a given ϕ we associate the vector $\mu(\phi) = (\mu_1(\phi), \dots, \mu_I(\phi))$, whose coordinates are

$$\mu_i(\phi) \doteq \sum_j \phi_{ij} \mu_{ij}, \quad i \in I; \quad (8)$$

this is the vector of mean long-run service rates provided to the types $i \in I$, if each server j allocates its time to serving type i at the average rate ϕ_{ij} .

Consider also a different vector-function of a matrix ϕ . Namely, let the vector $\rho(\phi) = (\rho_1(\phi), \dots, \rho_J(\phi)) \in R^J$ be defined as follows:

$$\rho_j(\phi) \doteq \sum_i \phi_{ij} \quad \forall j \in J. \quad (9)$$

Each component $\rho_j(\phi)$ is naturally interpreted as the total “utilization” of server j , given the average rates at which its time is allocated to service of different types i are given by ϕ_{ij} . (Again, we do not assume a priori that $\rho_j(\phi) \leq 1$.)

Definition 1 We define \mathcal{M} to be the set of $\mu(\phi)$ corresponding to all possible ϕ , satisfying the condition

$$\rho(\phi) \leq \mathbf{1} . \quad (10)$$

Further let \mathcal{M}^* denote the set of all maximal elements $\mu \in \mathcal{M}$ such that $\mu \in R_{++}^I$. ($\mu \in \mathcal{M}$ is maximal if $\mu \leq \zeta \in \mathcal{M}$ implies $\zeta = \mu$.)

Note that \mathcal{M} is a bounded convex polyhedron in R_+^I . We assume that \mathcal{M} is non-degenerate (i.e., has dimension I), which is equivalent to assuming that each customer type i can be served at non-zero rate μ_{ij} by at least one server j . The set \mathcal{M} is in fact the closure of our system's *stability region* \mathcal{M}^0 , which is the set of arrival rate vectors $\lambda = (\lambda_1, \dots, \lambda_I)$ such that $\lambda < \mu(\phi)$ for some ϕ satisfying (10). (Cf. [15, 13, 6, 2, 1, 14].)

Definition 2 We say that the condition of First Order Complete Resource Pooling (FO-CRP) holds for a vector λ , if λ lies within the interior of one of the $((I-1)$ -dimensional) outer faces of \mathcal{M} and $\lambda \in \mathcal{M}^*$. If, in addition, the matrix ϕ such that $\lambda = \mu(\phi)$ and (10) hold is unique, then we say that Complete Resource Pooling (CRP) condition holds.

Remark. The CRP condition given above is the same as in [12], and it is equivalent to that introduced earlier in [9, 20] (see Assumption 3.4, Theorem 5.3 and Corollary 5.4 in [20] for a summary). The fact of equivalence will, in particular, follow from the results of this section.

When the FO-CRP condition holds, let us denote by $\nu = (\nu_1, \dots, \nu_I)$ the (unique up to a scaling) “outer” normal vector to the polyhedron \mathcal{M} at the point λ . Note, that $\nu \in R_{++}^I$. (Otherwise, if some $\nu_i \leq 0$, a small increase of the component λ_i would produce a vector $\lambda' \geq \lambda$, $\lambda' \neq \lambda$, and such that $\lambda' \in \mathcal{M}$ - a contradiction to the maximality of λ .) For concreteness we use the normal vector ν^* , which is the vector defined uniquely by the additional requirement that $\|\nu^*\| = 1$. The components ν_i^* of the vector ν^* are sometimes called the *workload contributions* of customers of the different types i (see [9, 20, 12]); in this paper will call them *customer workload contributions*, to make a distinction from the server workload contributions introduced in Section 6.

The FO-CRP condition for λ implies, in particular, that

$$\nu^* \cdot \lambda = \max_{\mu \in \mathcal{M}} \nu^* \cdot \mu = \max_{\phi: \rho(\phi) \leq \mathbf{1}} \nu^* \cdot \mu(\phi) ; \quad (11)$$

this in turn implies that, for any matrix ϕ such that (10) holds and $\lambda = \mu(\phi)$, in fact, the equality in (10) must hold, namely, we have

$$\lambda = \mu(\phi) \text{ and } \rho(\phi) = 1 . \quad (12)$$

Given λ satisfying FO-CRP condition, for each $j \in J$, let us denote

$$I_j = \arg \max_i \nu_i^* \mu_{ij} \doteq \{i \in I \mid \nu_i^* \mu_{ij} = \max_l \nu_l^* \mu_{lj}\}.$$

Any pair (ij) such that $i \in I_j$ is called *basic activity*; therefore I_j indicates the set of basic activities for server j . It is easy to see from (11), that, for any ϕ satisfying (12), $\phi_{ij} > 0$ implies $i \in I_j$.

Lemma 1 *If λ satisfies the FO-CRP condition, then the corresponding graph \mathcal{G}^* with nodes being types i and servers j , and arcs being basic activities, is connected.*

Proof. Suppose not. Consider any breakdown of the graph \mathcal{G}^* into two components, $\mathcal{G}^{(1)}$ and $\mathcal{G}^{(2)}$, disconnected from each other. For $m = 1, 2$, denote by $I^{(m)}$ and $J^{(m)}$ the set of types and servers, respectively, within the component $\mathcal{G}^{(m)}$. By our construction, for any $m = 1, 2$, $j \in J^{(m)}$ implies $I_j \subseteq I^{(m)}$.

Consider any ϕ satisfying (12). Recall that $\phi_{ij} > 0$ implies $i \in I_j$. Let us fix a small $\delta > 0$, and consider vector ν^{**} which is obtained from ν^* by the following modification: $\nu_i^{**} = \nu_i^*(1 + \delta)$ if $i \in I^{(1)}$, and $\nu_i^{**} = \nu_i^*$ if $i \in I^{(2)}$. Since there is no arc connecting $\mathcal{G}^{(1)}$ and $\mathcal{G}^{(2)}$, if δ is small enough, then ϕ solves the problem

$$\max_{\phi: \rho(\phi) \leq 1} \nu^{**} \cdot \mu(\phi)$$

as well as the problem in the RHS of (11). In other words, $\lambda = \mu(\phi)$ solves the problem $\max_{\mu \in \mathcal{M}} \nu^{**} \cdot \mu$, as well as $\max_{\mu \in \mathcal{M}} \nu^* \cdot \mu$. This means that ν^{**} is a normal (different from ν^*) to the boundary of \mathcal{M} at point λ - a contradiction to the FO-CRP condition. ■

Now, with *any* matrix ϕ let us associate graph $\mathcal{G}(\phi)$ with nodes being types i and servers j , and arcs (ij) corresponding to pairs (ij) with $\phi_{ij} > 0$.

Lemma 2 (i) *If FO-CRP holds, then there exists ϕ satisfying (12) and such that $\phi_{ij} > 0$ if and only if $i \in I_j$.*

(ii) *The FO-CRP condition for λ holds if and only if $\lambda \in \mathcal{M}^*$ and there exists ϕ such that (12) holds and the graph $\mathcal{G}(\phi)$ is connected.*

(iii) *If CRP condition holds, then ϕ satisfying (12) is unique, the graph $\mathcal{G}(\phi) = \mathcal{G}^*$ (that is, $\phi_{ij} > 0$ if and only if $i \in I_j$), and this graph is a tree.*

Proof. (i) Consider arbitrary ϕ' satisfying $\rho(\phi') = 1$, and such that $\phi'_{ij} > 0$ if and only if $i \in I_j$. Note that $\nu^* \cdot \mu(\phi') = \nu^* \cdot \lambda$, because the condition on ϕ' guarantees that ϕ' solves the problem in the RHS of (11). Let

$$\lambda'' = \frac{\lambda - \mu(\delta\phi')}{1 - \delta},$$

where $0 < \delta < 1$ is fixed. We have

$$\nu^* \cdot \lambda'' = \frac{\nu^* \cdot \lambda - \delta \nu^* \cdot \mu(\phi')}{1 - \delta} = \nu^* \cdot \lambda,$$

and we can always choose δ to be small enough, so that λ'' lies in the interior of the same face of \mathcal{M} as λ does. Then, there exists ϕ'' such that $\lambda'' = \mu(\phi'')$, $\rho(\phi'') = 1$, and $\phi''_{ij} > 0$ implies $i \in I_j$. It is easy to verify directly that $\phi = (1 - \delta)\phi'' + \delta\phi'$ is a matrix with the properties we seek.

(ii) Necessity follows from (i). Let us prove sufficiency. Let ϕ be a matrix such that (12) holds, $\lambda \in \mathcal{M}^*$, and the graph $\mathcal{G}(\phi)$ is connected. Since $\lambda \in \mathcal{M}^*$, there exists an outer normal vector ν^* to the boundary of \mathcal{M} at point λ , and it is such that $\nu^* \neq 0$, $\nu^* \in R_+^I$. Consequently, ϕ solves the problem (in the RHS of) (11). From this we conclude that $\nu^* \in R_{++}^I$ - otherwise (if $\nu_i^* = 0$ for some i), ϕ could not be a solution to (11), since $\mathcal{G}(\phi)$ is connected. Finally, the normal ν^* must be unique (up to scaling). Indeed, consider any other vector $\nu^{**} \in R_{++}^I$, which is not a scaled version of ν^* , that is, $\max_i \nu_i^{**}/\nu_i^* > \min_i \nu_i^{**}/\nu_i^*$. Then, connectedness of $\mathcal{G}(\phi)$ easily implies that, since ϕ solves (11), it cannot possibly solve (11) with ν^* replaced by ν^{**} . Therefore, ν^{**} cannot be a normal to \mathcal{M} at point λ .

(iii) The definition of CRP and statements (i) and (ii) of the lemma immediately apply the uniqueness of ϕ satisfying (12), the fact that $\mathcal{G}(\phi) = \mathcal{G}^*$ and that this graph is connected. It remains to show that $\mathcal{G}(\phi)$ must be a tree. Suppose not. Let us pick any cycle on this graph. It is easy to see that we can “perturb” the (strictly positive) elements ϕ_{ij} along the arcs of the cycle, so as to produce a matrix $\phi' \neq \phi$, such that $\mu(\phi') = \mu(\phi) = \lambda$ and $\rho(\phi') = 1$, a contradiction to the uniqueness of ϕ . ■

Remark. Just as in the case of $\text{Gc}\mu$ scheduling rule, studied for an input-queued model in [12], we emphasize here that the notion of a basic activity is *not* utilized in any way (neither explicit nor implicit) by the MinDrift routing algorithm. (The algorithm need not know which activities are basic.) It is only used as a tool for the analysis of the algorithm. Similarly, the algorithm need not know the values of workload contributions.

6 Equivalent Characterization of CRP Condition in Terms of the Output-Queued Model

In this section we give an equivalent (“dual”) characterization of the CRP condition, and introduce notions and results which will be used in the analysis of our OQ-model.

First, let us give a somewhat different (although closely related) interpretation of the matrix ϕ and functions $\mu(\phi)$ and $\rho(\phi)$, defined in Section 5. Suppose a matrix ϕ is given, and assume that customers of a type i arrive (routed) to a server j at the average rate $\mu_{ij}\phi_{ij}$. Then, $\mu_i(\phi)$, $i \in I$, is the total average rate at which type i customers arrive in the system, and

$$\rho_j(\phi) = \sum_i \phi_{ij} \equiv \sum_i (\mu_{ij}\phi_{ij})\mu_{ij}^{-1}, \quad j \in J,$$

is the average rate at which the *work* (i.e., the required amount of processing time) arrives (routed) to server j . (We used the convention that $(\mu_{ij}\phi_{ij})\mu_{ij}^{-1} = 0$ if $\mu_{ij} = 0$.)

Definition 3 We define the server utilization region $\mathcal{K} \subseteq R_+^J$ to be the set of all possible values of $\rho(\phi)$ with ϕ satisfying condition

$$\mu(\phi) \geq \lambda. \quad (13)$$

Further let \mathcal{K}_* denote the set of all minimal elements $\rho \in \mathcal{K}$ such that $\rho \in R_{++}^J$. ($\rho \in \mathcal{K}$ is minimal if $\rho \geq \zeta \in \mathcal{K}$ implies $\zeta = \rho$.)

Region \mathcal{K} is a convex polyhedron in R_+^J , and it is non-degenerate (i.e., has dimension J) as long as \mathcal{M} is non-degenerate. Note that \mathcal{K} is unbounded, but it is of course “bounded below”, say by 0, since it lies in the positive orthant.

Lemma 3 (i) The FO-CRP condition for a fixed vector λ holds if and only if the following is true:

- (a) vector $\mathbf{1} \in R^J$ lies within the interior of one of the $((J-1)$ -dimensional faces of \mathcal{K} ,
- (b) $\mathbf{1} \in \mathcal{K}_*$.

(ii) When FO-CRP condition for λ (or, equivalently, (i)(a) and (i)(b)) holds, then the unique (up to a scaling by positive constant) outer normal vector $-\alpha^*$ to the polyhedron \mathcal{K} at the point $\mathbf{1}$ is such that $\alpha^* = (\alpha_1^*, \dots, \alpha_J^*) \in R_{++}^J$, and α^* is related to the vector ν^* as follows:

$$\alpha_j^* = \max_i \mu_{ij} \nu_i^*, \quad j \in J, \quad (14)$$

$$\nu_i^* = \min_j \alpha_j^* / \mu_{ij}, \quad i \in I. \quad (15)$$

In addition:

- (c) $i \in I_j$ (that is, activity (ij) is basic) if and only if $j \in J_i$, where

$$J_i \doteq \arg \min_j \alpha_j^* / \mu_{ij}; \quad (16)$$

(d) any matrix ϕ satisfying $\rho(\phi) = \mathbf{1}$ and (13), in fact, satisfies (12);

(e) we have

$$\alpha^* \cdot \mathbf{1} = \nu^* \cdot \lambda. \quad (17)$$

(iii) The CRP condition for a fixed vector λ holds if and only if (i)(a), (i)(b) and the following property hold:

(f) the matrix ϕ satisfying $\rho(\phi) = \mathbf{1}$ and (13) is unique.

When CRP does hold, the matrix ϕ is in fact the unique solution of (12).

Proof. (i) Let us prove the necessity of (a) and (b). Consider the vector α^* defined by (14). (Note that $\alpha^* \in R_{++}^J$.) Then (15) holds. Indeed, for a fixed i , we have $\alpha_j^* = \mu_{ij} \nu_i^*$ if (ij) is basic, and $\alpha_j^* > \mu_{ij} \nu_i^*$ otherwise. (Incidentally, this means that $i \in I_j$ is equivalent to $j \in J_i$.)

Let choose any matrix ϕ which solves (12) and such that $\mathcal{G}(\phi) = \mathcal{G}^*$. (Recall that graph \mathcal{G}^* is connected.) Then, since this ϕ is such that $\phi_{ij} > 0$ implies $j \in J_i$, it is easy to observe that ϕ solves the problem

$$\min_{\phi: \mu(\phi) \geq \lambda} \alpha^* \cdot \rho(\phi), \quad (18)$$

or, equivalently, $\mathbf{1}$ solves the problem

$$\min_{\rho \in \mathcal{K}} \alpha^* \cdot \rho. \quad (19)$$

(Compare problems (18) and (19) to problem (11).) This means, in particular, that $\mathbf{1}$ is a minimal element of \mathcal{K} , and α^* is a normal to the region \mathcal{K} at point $\mathbf{1}$. Since $\mathcal{G}(\phi)$ is connected, it is easy to see from (18) that $\rho = \mathbf{1}$ could not possibly solve the problem (19) with α^* replaced by any other non-zero vector $\alpha^{**} \in R_+^J$, unless $\alpha^{**} = c\alpha^*$ for some $c > 0$. This completes the proof of necessity of (a) and (b).

The sufficiency of (a) and (b) follows simply by the symmetry between the definitions of FO-CRP condition (for λ) and condition (a)-(b) (expressed in terms of vector $\mathbf{1}$). Namely, if for the vector $-\mathbf{1}$ and the region $-\mathcal{K}$ we define a condition analogous to FO-CRP for λ and region \mathcal{M} , this will be exactly the condition (a)-(b). Thus, from this condition ((a)-(b)) we can obtain a condition analogous to (a)-(b), but with $\mathbf{1}$ and \mathcal{K} replaced by $-\lambda$ and $-\mathcal{M}$, respectively; this latter condition is exactly our original FO-CRP.

(ii) As part of the proof of (i), we already proved all the properties stated in (ii), except (d) and (e). If (d) would not hold, then λ could not be a maximal element of \mathcal{M} . Property (e) follows from the fact that any matrix ϕ , chosen as in the proof of (i), satisfies (12) and solves both problems (11) and (18).

(iii) This follows from (i), (ii) and the definition of CRP. ■

When FO-CRP condition holds, the components α_j^* of the vector α^* we will call *server workload contributions* of different servers j .

7 The Heavy Traffic Regime

In this section we introduce the notion of a *sequence of queueing systems in heavy traffic*. Suppose a vector λ satisfying the CRP condition is fixed. Associated with this λ are the unique matrix ϕ satisfying (12), and the corresponding normal vectors $\nu^* \in R^I$ and $\alpha^* \in R^J$. (We remark, that all the definitions and facts in this section are valid when the weaker FO-CRP condition holds, with *arbitrary* fixed ϕ satisfying (12). However, for our main results in Section 8 the stronger CRP condition is essential.)

The quantity

$$X(t) \doteq \sum_{i=1}^I \nu_i^* Q_i(t), \quad t \geq 0 ,$$

we will call *customer workload* of the system. The customer workload process $X(\cdot)$ is of the primary interest in the analysis of the input-queued model [8, 9, 20, 3, 12].

For the output-queued model, we define a different (although closely related, in the sense specified later) notion of *server workload*:

$$^u X(t) \doteq \sum_{j=1}^J \alpha_j^* U_j(t), \quad t \geq 0.$$

In addition, we define *Q-estimated server workload* as

$$^q X(t) \doteq \sum_{j=1}^J \alpha_j^* {}^q U_j(t), \quad t \geq 0.$$

Informally speaking, for a service discipline satisfying constraints (d0)-(d2), ${}^q X(t)$ is a “good” (asymptotically exact) estimate of the server workload ${}^u X(t)$.

Since for any pair of $i \in I$ and $j \in J$ the inequality $\alpha_j^*/\mu_{ij} \geq \nu_i^*$ holds, we observe that the *Q*-estimated server workload cannot be less than customer workload:

$${}^q X(t) \equiv \sum_{i=1}^I \sum_{j=1}^J (\alpha_j^*/\mu_{ij}) Q_{ij} \geq X(t) . \quad (20)$$

We also have the following inequality, which we record for future reference:

$${}^q X^r(t) \leq C_0 X^r(t), \quad t \geq 0, \quad (21)$$

with

$$C_0 = \max_{(ij): \mu_{ij} > 0} \frac{\alpha_j^*/\mu_{ij}}{\nu_i^*} . \quad (22)$$

We now consider a sequence of queueing systems, indexed by $r \in \mathcal{R} = \{r_1, r_2, \dots\}$, where $r_n > 0$ for all n and $r_n \uparrow \infty$ as $n \rightarrow \infty$. (Hereafter in this paper, “ $r \rightarrow \infty$ ” means that r goes to infinity along the sequence \mathcal{R} , or some subsequence of \mathcal{R} ; the choice of the subsequence will be either explicit or clear from the context.) Each system $r \in \mathcal{R}$ has, as before, I customer types and J servers. The primitives and the processes corresponding to a system $r \in \mathcal{R}$ will be appended with a superscript r .

Assume that, for each type i , the mean arrival rate $\lambda_i^r = 1/E[u_i^r(1)]$ is such that

$$r(\lambda_i^r - \lambda_i) \rightarrow b_i , \quad r \rightarrow \infty , \quad (23)$$

where $b_i \in R$ is a fixed constant. Assume also convergence of the variance, that is

$$[\alpha_i^r]^2 \rightarrow \alpha_i^2, \quad r \rightarrow \infty. \quad (24)$$

In addition, we make the following technical assumption, needed, in particular, to apply Bramson's weak law estimates [4] (and establish (75) later on): uniformly over i and r ,

$$E[(u_i^r(1))^2 \mathbf{1}\{u_i^r(1) > x\}] \leq \eta(x), \quad x \geq 0, \quad (25)$$

where $\eta(\cdot)$, is a fixed function, $\eta(x) \rightarrow 0$ as $x \rightarrow \infty$.

For the initial interarrival times we assume that, for each i ,

$$u_i^r(0)/r \rightarrow 0, \quad r \rightarrow \infty.$$

Let $F_i^r(t)$, $t \geq 0$, denote the number of type i customers arrived in the system by time t , excluding "initial" customers present at time 0. Assumptions (23), (24) and (25) imply a functional central limit theorem (FCLT) for these arrival processes:

$$\{r^{-1}(F_i^r(r^2t) - \lambda_i^r r^2 t), t \geq 0\} \xrightarrow{w} \{\sigma_i B(t), t \geq 0\}, \quad (26)$$

where $\sigma_i^2 = \lambda_i^3 \alpha_i^2$, $B(\cdot)$ is a standard (zero drift, unit variance) Brownian motion, and \xrightarrow{w} denotes convergence in distribution (for processes in the standard Skorohod space of RCLL functions).

The service time distributions do *not* change with the parameter r . (This in particular means that the condition analogous to (25) trivially holds for the service times $v_{i,j}^r(1)$, uniformly on (i, j) and r .)

Let us denote by

$${}^{\Sigma}V_{ij}^r(l) \doteq \sum_{m=1}^l v_{ij}^r(m), \quad l = 0, 1, 2, \dots,$$

the total amount of work (i.e., total service time) brought to server j by the first l (newly arriving) type i customers routed to it. We extend the domain of ${}^{\Sigma}V_{ij}^r(\cdot)$ to all real non-negative $t \geq 0$ by adopting the convention that ${}^{\Sigma}V_{ij}^r(t) = {}^{\Sigma}V_{ij}^r(\lfloor t \rfloor)$. (We will use same domain extension convention throughout the paper, for other functions, which are originally defined on the integers, as well.) For ${}^{\Sigma}V_{ij}^r$ we have the following FCLT:

$$\{r^{-1}({}^{\Sigma}V_{ij}^r(r^2t) - \mu_{ij}^{-1} r^2 t), t \geq 0\} \xrightarrow{w} \{\beta_{ij} B(t), t \geq 0\}. \quad (27)$$

For each $i \in I$, let us fix a set of integer valued non-decreasing non-negative functions $({}^s N_{ij}(l), l = 0, 1, 2, \dots)$, $j \in J$, satisfying the following conditions:

$$\sum_j {}^s N_{ij}(l) = l, \quad l = 0, 1, 2, \dots, \quad (28)$$

$$\max_{l \geq 0} |{}^s N_{ij}(l) - \frac{\mu_{ij}\phi_{ij}}{\lambda_i} l| < \infty, \quad j \in J, \quad (29)$$

$${}^s N_{ij}(l) \equiv 0 \text{ for } l = 0, 1, 2, \dots, \quad j \notin J_i. \quad (30)$$

The value of ${}^s N_{ij}(l)$ is interpreted as the number of type i customers routed to server j , out of the first l type i customers arrived in the system. Then, it is clear that, for each flow i , the functions ${}^s N_{ij}(\cdot)$ define a fixed (“static”) pattern of routing customers to the servers in J_i , such that, for any l , the fractions of customers routed to different servers $j \in J_i$ closely track the values $\mu_{ij}\phi_{ij}/\lambda_i$; recall that $\sum_j \mu_{ij}\phi_{ij}/\lambda_i = 1$. (The MinDrift rule does not require any knowledge of this static routing pattern - it is only used as a tool for the analysis!) Such functions can be, for example, constructed recursively as follows. We set ${}^s N_{ij}(0) = 0$ for all $j \in J_i$. For each $l = 1, 2, \dots$, we set ${}^s N_{ij}(l) = {}^s N_{ij}(l-1) + 1$ for one of the $j \in J_i$ with the smallest value of ${}^s N_{ij}(l-1) - (\mu_{ij}\phi_{ij}/\lambda_i)l$, and ${}^s N_{ij}(l) = {}^s N_{ij}(l-1)$ for all other j . (The “ties” between j are broken arbitrarily, for example, in favor of the smallest one.)

For $i \in I$, let us denote by

$$A_i^r(t) \doteq \sum_{j \in J} \alpha_j^{*\Sigma} V_{ij}^r({}^s N_{ij}(F_i^r(t))) \equiv \sum_{j \in J} \alpha_j^* \sum_{m=1}^{{}^s N_{ij}(F_i^r(t))} v_{ij}^r(m), \quad t \geq 0,$$

the total amount of server workload brought to the system by the new arrivals of flow i by time $t \geq 0$ *assuming the arrivals would be routed according to the (fixed) functions ${}^s N_{ij}(\cdot)$* . From (26)-(30) we obtain the following FCLT for the sequence of processes $A_i^r(\cdot)$:

$$\{r^{-1}(A_i^r(r^2t) - \lambda_i^r \frac{\sum_j \phi_{ij}\alpha_j^*}{\lambda_i} r^2 t), \quad t \geq 0\} \xrightarrow{w} \{{}^u \sigma_i B(t), \quad t \geq 0\}, \quad (31)$$

where

$$\begin{aligned} {}^u \sigma_i^2 &\doteq \lambda_i [\sum_j \phi_{ij}\alpha_j^*]^2 \alpha_i^2 + \sum_j \phi_{ij}\mu_{ij}\beta_{ij}^2 [\alpha_j^*]^2 \\ &= [\nu_i^*]^2 [\lambda_i^3 \alpha_i^2 + \sum_j \phi_{ij}\mu_{ij}^3 \beta_{ij}^2]. \end{aligned} \quad (32)$$

From (31) and (17) we obtain the following FCLT for the sequence of processes $\sum_i A_i^r(\cdot)$:

$$\{r^{-1}(\sum_i A_i^r(r^2t) - [\sum_j \alpha_j^*] r^2 t), \quad t \geq 0\} \xrightarrow{w} \{at + \sigma B(t), \quad t \geq 0\}, \quad (33)$$

where

$$\sigma^2 \doteq \sum_i {}^u \sigma_i^2, \quad a \doteq \nu^* \cdot b. \quad (34)$$

8 Main Results

For each value of the (scaling) parameter $r \in \mathcal{R}$, let us consider the following processes. Let $U^r(\cdot)$ and ${}^q U^r(\cdot)$ be the (vector) processes, representing (unfinished) work and Q -estimated (unfinished) work, respectively, at different servers; let ${}^u X^r(\cdot)$, ${}^q X^r(\cdot)$, and $X^r(\cdot)$ denote the scalar processes representing server workload, Q -estimated server workload, and customer workload, respectively.

Assume that, in a system with index $r \in \mathcal{R}$, each server j , at any time t , incurs a *holding cost* at the (instantaneous) rate of

$$C_j^r(U_j^r(t)) = C_j(U_j^r(t)/r) ;$$

where $C_j(\cdot)$ are fixed convex increasing functions, with the additional properties described in Section 4.

Note that in our asymptotic regime the cost function is “rescaled” as the parameter r changes. (In other words, in a system with index r , the holding cost rate corresponding to the unfinished work $U_j^r(t)$ is $C_j(U_j^r(t)/r)$ instead of $C_j(U_j^r(t))$.) Notice, however, that in the special case (described in Section 4.4) when $C_j(\zeta) = \gamma_j \zeta^{\eta+1}$, with some fixed $\eta > 0$ and $\gamma_j > 0$, *the form of the corresponding MinDrift rule does not change with r* . Indeed, *in this case*, replacing in (2) $C'_j(U_j^r(t))$ with $C'_j(U_j^r(t)/r)/r$ simply does not change the routing rule.

For our main results, we need the notion of a fixed point. A vector ${}^o u \in R_+^J$ will be called a *fixed point* if,

$$[C'_1({}^o u_1), \dots, C'_J({}^o u_J)] = c \alpha^* , \quad (35)$$

for some constant $c \geq 0$. If we recall that each derivative $C'_j(\cdot)$ is continuous strictly increasing with $C'_j(0) = 0$, one deduces the following:

A fixed point ${}^o u$ corresponding to each $c \geq 0$ exists and is unique. Moreover, ${}^o u = 0$ for $c = 0$, and ${}^o u \in R_{++}^J$ (i.e. has all components strictly positive) for any $c > 0$.

Thus, the set of fixed points forms a one-dimensional manifold, which can be parameterized for example by the corresponding server workload values $\alpha^* \cdot {}^o u$. In addition, it is easy to verify the following property:

A fixed point ${}^o u$ is the unique vector that minimizes $\sum_j C_j(u_j)$ among all vectors $u \in R_+^J$ with the same server workload, i.e. satisfying condition $\alpha^ \cdot u = \alpha^* \cdot {}^o u$.*

Indeed, if ${}^o u = 0$, the property is trivial. If ${}^o u \in R_{++}^J$, condition (35) implies that the (Lagrangian) function

$$\sum_j C_j(u_j) - c[\alpha^* \cdot u - \alpha^* \cdot {}^o u]$$

has zero gradient (with respect to u) at point ${}^o u$. Since this Lagrangian is strictly convex in R_+^J , it is minimized by ${}^o u$. Then, the desired property follows from the Kuhn-Tucker theorem.

Let us define the *diffusion scaling* operator $\tilde{\Gamma}^r$, which acts on a scalar function $\Xi = (\Xi(t), t \geq 0)$ as follows:

$$(\tilde{\Gamma}^r \Xi)(t) \doteq \frac{1}{r} \Xi(r^2 t) , \quad (36)$$

and is applied to vector-functions componentwise.

Let us consider the following *diffusion-scaled* processes: $\tilde{u}^r = \tilde{\Gamma}^r U^r$, ${}^q\tilde{u}^r = \tilde{\Gamma}^r {}^q U^r$, ${}^u\tilde{x}^r = \tilde{\Gamma}^r {}^u X^r$, ${}^q\tilde{x}^r = \tilde{\Gamma}^r {}^q X^r$, $\tilde{x}^r = \tilde{\Gamma}^r X^r$.

8.1 Optimality of MinDrift(U) Rule.

Assume that the initial (scaled) amounts of unfinished work are deterministic and converging:

$$\tilde{u}^r(0) \rightarrow \tilde{u}(0) , \quad (37)$$

where $\tilde{u}(0)$ is a fixed point, as defined above. Consequently, ${}^u\tilde{x}^r(0) = \alpha^* \cdot \tilde{u}^r(0) \rightarrow \alpha^* \cdot \tilde{u}(0) \doteq \tilde{w}(0)$.

Let us define the following one-dimensional reflected Brownian motion $\tilde{x} = \{\tilde{x}(t), t \geq 0\}$:

$$\tilde{x}(t) = \tilde{w}(0) + at + \sigma B(t) + \tilde{y}(t) , \quad (38)$$

where $B(\cdot)$ is a standard Brownian motion,

$$\tilde{y}(t) \doteq -[0 \wedge \inf_{0 \leq \xi \leq t} \{\tilde{w}(0) + a\xi + \sigma B(\xi)\}] , \quad (39)$$

and the drift a and diffusion coefficient σ are given in (34) and (32).

Theorem 1 Consider the sequence of queueing systems in heavy traffic, as introduced in Section 7. Assume initial condition (37). Let \tilde{x} be a reflected Brownian motion defined by (38)-(39).

(i) Suppose that the service discipline is such that the routing rule is MinDrift(U) with cost functions $C_i^r(\cdot)$, for each value of the parameter r , and each server employs an arbitrary work-conserving scheduling rule (namely, the server is not allowed to idle when there is unfinished work in its queue). Then, as $r \rightarrow \infty$,

$${}^u\tilde{x}^r \xrightarrow{w} \tilde{x} ,$$

and

$$\tilde{u}^r \xrightarrow{w} \tilde{u} ,$$

where, for each $t \geq 0$, the vector $\tilde{u}(t)$ is the fixed point that is (uniquely) determined by $\alpha^* \cdot \tilde{u}(t) = \tilde{x}(t)$.

(ii) The service discipline defined in (i) is asymptotically optimal within the class of service disciplines satisfying condition (d0) in that it minimizes the server workload and the unfinished work cost rate at all times. More precisely, let \tilde{u}_G^r and ${}^u\tilde{x}_G^r$ be the scaled unfinished work and server workload processes corresponding to an arbitrary service discipline G (and appropriately constructed on a common probability space with the sequence of processes defined in (i)). Then, with probability 1, for any time $t \geq 0$,

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [{}^u\tilde{x}_G^r(\xi) - \tilde{x}(\xi)] \geq 0 \quad (40)$$

and

$$\liminf_{r \rightarrow \infty} \sum_j C_j(\tilde{u}_{j,G}^r(t)) \geq \sum_j C_j(\tilde{u}_j(t)) . \quad (41)$$

As a corollary, with probability 1, for any $T > 0$,

$$\liminf_{r \rightarrow \infty} \int_0^T \sum_j C_j(\tilde{u}_{j,G}^r(t)) dt \geq \lim_{r \rightarrow \infty} \int_0^T \sum_j C_j(\tilde{u}_j^r(t)) dt = \int_0^T \sum_j C_j(\tilde{u}_j(t)) dt . \quad (42)$$

The proof of Theorem 1 is the subject of Sections 9 and 10.

8.2 Optimality of MinDrift(Q) Rule.

Assume that for each r the initial state of the system at time 0 is deterministic, and it conforms to the conditions (d1) and (d2) on a service discipline (which are assumed in Theorem 2 below). In particular, for each pair of i and j , server j has in its queue at most one customer of type i whose service has already started, and the realizations of service times of the customers whose service has not yet started are not known to the server. For the initial residual service times $v_{i,j}^r(0)$ (if any) of the customers whose service has already started, we assume

$$v_{i,j}^r(0)/r \rightarrow 0, \quad r \rightarrow \infty.$$

Finally, assume that the initial (scaled) amounts of Q -estimated unfinished work are converging:

$${}^q\tilde{u}^r(0) \rightarrow \tilde{u}(0) , \quad (43)$$

where $\tilde{u}(0)$ is a fixed point, as defined above.

It follows from the above initial conditions that

$${}^q\tilde{x}^r(0) = \alpha^* \cdot {}^q\tilde{u}^r(0) \rightarrow \alpha^* \cdot \tilde{u}(0) \doteq \tilde{w}(0) \quad (44)$$

and, in addition, with probability 1,

$$\tilde{u}^r(0) \rightarrow \tilde{u}(0) \text{ and } {}^u\tilde{x}^r(0) \rightarrow \tilde{w}(0) . \quad (45)$$

For the fixed $\tilde{w}(0)$, we consider a one-dimensional reflected Brownian motion $\tilde{x} = \{\tilde{x}(t), t \geq 0\}$, defined in (38).

Theorem 2 Consider the sequence of queueing systems in heavy traffic, as introduced in Section 7, and with the initial conditions described above in Section 8.2.

(i) Suppose that the service discipline is such that the routing rule is $\text{MinDrift}(Q)$ with cost functions $C_j^r(\cdot)$, for each value of the parameter r , and each server employs an arbitrary work-conserving scheduling rule satisfying conditions (d1)-(d2). Then, as $r \rightarrow \infty$,

$${}^q\tilde{x}^r \xrightarrow{w} \tilde{x}, \quad {}^u\tilde{x}^r \xrightarrow{w} \tilde{x},$$

and

$${}^q\tilde{u}^r \xrightarrow{w} \tilde{u}, \quad \tilde{u}^r \xrightarrow{w} \tilde{u},$$

where, for each $t \geq 0$, the vector $\tilde{u}(t)$ is the fixed point that is (uniquely) determined by $\alpha^* \cdot \tilde{u}(t) = \tilde{x}(t)$.

(ii) The service discipline defined in (i) is asymptotically optimal within the class of service disciplines satisfying conditions (d0)-(d2) in that it minimizes the both the server workload and the Q -estimated server workload, and the unfinished work cost rate at all times. More precisely, let \tilde{u}_G^r , ${}^q\tilde{u}_G^r$, ${}^u\tilde{x}_G^r$ and ${}^q\tilde{x}_G^r$ be the scaled unfinished work, Q -estimated unfinished work, server workload, and Q -estimated server workload processes, respectively, corresponding to an arbitrary service discipline G satisfying (d0)-(d2) (and appropriately constructed on a common probability space with the sequence of processes in (i)). Then, with probability 1, for any time $t \geq 0$,

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [{}^q\tilde{x}_G^r(\xi) - \tilde{x}(\xi)] = \liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [{}^u\tilde{x}_G^r(\xi) - \tilde{x}(\xi)] \geq 0 \quad (46)$$

and

$$\liminf_{r \rightarrow \infty} \sum_j C_j({}^q\tilde{u}_{j,G}^r(t)) = \liminf_{r \rightarrow \infty} \sum_j C_j(\tilde{u}_{j,G}^r(t)) \geq \sum_j C_j(\tilde{u}_j(t)). \quad (47)$$

As a corollary, with probability 1, for any $T > 0$,

$$\lim_{r \rightarrow \infty} \int_0^T \sum_j C_j({}^q\tilde{u}_j^r(t)) dt = \lim_{r \rightarrow \infty} \int_0^T \sum_j C_j(\tilde{u}_j^r(t)) dt = \int_0^T \sum_j C_j(\tilde{u}_j(t)) dt \quad (48)$$

and

$$\liminf_{r \rightarrow \infty} \int_0^T \sum_j C_j({}^q\tilde{u}_{j,G}^r(t)) dt = \liminf_{r \rightarrow \infty} \int_0^T \sum_j C_j(\tilde{u}_{j,G}^r(t)) dt \geq \int_0^T \sum_j C_j(\tilde{u}_j(t)) dt. \quad (49)$$

The proof of Theorem 2 is essentially a slightly modified (and extended) version of that of Theorem 1. It is outlined in Section 11.

8.3 Customer Workload Minimization under MinDrift(Q) Rule.

Suppose we are within the conditions of Theorem 2. For the initial customer workloads $\tilde{x}^r(0)$ we “automatically” (by (20) and (44)) have

$$\limsup_r \tilde{x}^r(0) \leq \lim_r {}^q\tilde{x}^r(0) = \tilde{w}(0) .$$

Suppose, in addition, that, in fact, $\tilde{x}^r(0)$ converges to the same limit as ${}^q\tilde{u}^r(0)$ does:

$$\lim_r \tilde{x}^r(0) = \lim_r {}^q\tilde{x}^r(0) = \tilde{w}(0) , \quad (50)$$

which is equivalent to the condition that $\lim_r \tilde{q}_{ij}^r(0) = 0$ for every non-basic activity (ij) .

Now, any service discipline in the OQ-system, satisfying conditions (d0)-(d2), is within the class of disciplines for the corresponding IQ-system studied in [12] (see Remark 2 in Section 3 of this paper). In particular, Theorem 1 in [12] establishes that an RBM with exactly same distribution as the RBM \tilde{x} (defined in this paper by (38)) is in fact the stochastic *lower* bound of any limit of the customer workload process \tilde{x}^r , under any service discipline satisfying conditions (d0)-(d2) (but not necessarily (IR)), which includes service disciplines for both OQ- and IQ-systems.

However, from (20) we have a pathwise relation

$$\tilde{x}^r(t) \leq {}^q\tilde{x}^r(t), \quad t \geq 0, \quad (51)$$

and, by Theorem 2(i), under MinDrift(Q) rule, ${}^q\tilde{x}^r \xrightarrow{w} \tilde{x}$. Thus, \tilde{x} is both the lower and upper (stochastic) bound of \tilde{x}^r and, therefore, $\tilde{x}^r \xrightarrow{w} \tilde{x}$. We have proved the following result, which basically says that MinDrift(Q) rule minimizes customer workload among virtually all service disciplines in either the OQ- or IQ-system.

Theorem 3 *Suppose, the conditions of Theorem 2 and, in addition, condition (50) hold.*

(i) For the service discipline described in Theorem 2(i), we in addition have

$$\tilde{x}^r \xrightarrow{w} \tilde{x}. \quad (52)$$

(ii) The service discipline described in Theorem 2(i) asymptotically minimizes customer workload among all service disciplines, satisfying conditions (d0)-(d2), in either OQ- or IQ-system. Namely, the customer workload process \tilde{x}_G^r under any such discipline G can be constructed on a common probability space with the RBM \tilde{x} , so that, with probability 1, for any time $t \geq 0$,

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [\tilde{x}_G^r(\xi) - \tilde{x}(\xi)] \geq 0 .$$

8.4 A Necessary Condition for Workload Minimization under Any Service Discipline: Vanishing Non-Basic Queues.

Theorems 2 and 3 show that, roughly speaking, the MinDrift(Q) rule minimizes both server and customer workload in the heavy traffic limit. The following Theorem 4 demonstrates that (either customer or server) workload in an OQ-system can be minimized by some discipline only if, under this discipline, the (scaled) queue lengths \tilde{q}_{ij}^r corresponding to non-basic activities (ij) vanish in the limit.

Theorem 4 *Suppose, the conditions of Theorem 2 and, in addition, condition (50) hold. Suppose, under some service discipline in OQ-system, satisfying conditions (d0)-(d2), either (52) or*

$${}^u\tilde{x}^r \xrightarrow{w} \tilde{x} \quad (53)$$

or

$${}^q\tilde{x}^r \xrightarrow{w} \tilde{x} \quad (54)$$

holds. Then, for any $t \geq 0$,

$$\tilde{q}_{ij}^r(t) \xrightarrow{P} 0 \quad \text{for any non-basic activity } (ij). \quad (55)$$

The proof is presented in Section 12.

9 Fluid Sample Paths under the MinDrift Rule

9.1 Fluid Sample Paths: Definition and Basic Properties.

In this section we study the sequence of processes introduced in the previous section under the *fluid* (or “law of large numbers”) scaling and under the MinDrift-rule. More precisely, we need to consider only *sample paths* of the processes under this scaling, and then their limits, which we formally define below and call *fluid sample paths* (FSP). The key property of FSPs that must be established (in Theorem 5 below) is that, as time increases to infinity, the queue length vector converges to a fixed point. This attraction property is used to prove (in the next section) the state space collapse property, i.e. the property that the limit of the sequence of *diffusion scaled* processes is a process “living” on the manifold of fixed points.

Throughout Section 9 we assume the CRP condition. However, all definitions and results of this section hold under the weaker FO-CRP condition, with *arbitrary* fixed ϕ satisfying (12) and $\mathcal{G}(\phi) = \mathcal{G}^*$; in other words, the FSP definition and key properties do *not* require a solution ϕ of (12) to be unique.

First, we introduce some additional (random) functions, associated with the process for each value of the scaling parameter r . (The functions $F_i^r(t)$, $Q_i^r(t)$, and $X^r(t)$, have already been defined earlier.)

Denote by $G_{ij}^r(t)$ the amount of time within $[0, t]$ that server j was serving type i customers.

For each pair (i, j) we define $N_{ij}^r(n)$ as the number of type i arrivals actually routed to server j , out of the first n new type i arrivals, and denote

$$H_{ij}^r(t) \doteq N_{ij}^r(F_i^r(t)) - {}^s N_{ij}^r(F_i^r(t)) .$$

Here, for any pair (i, j) , ${}^s N_{ij}^r(\cdot) \equiv {}^s N_{ij}(\cdot)$ for all r , where ${}^s N_{ij}(\cdot)$ are *non-random* functions fixed earlier and satisfying conditions (28)-(30).

We define (*server workload*) *regulation* process as follows:

$$Y^r(t) \doteq Y_{idle}^r(t) + Y_{route}^r(t) , \quad t \geq 0 ,$$

where

$$Y_{idle}^r(t) \doteq \sum_j \alpha_j^*(t - \sum_i G_{ij}^r(t)) ,$$

$$Y_{route}^r(t) \doteq \sum_i Y_{route,i}^r(t) ,$$

$$Y_{route,i}^r(t) \doteq \sum_j \alpha_j^* \mu_{ij}^{-1} H_{ij}^r(t) .$$

Function Y_{idle}^r is the regulation component due to physical idleness of the servers, Y_{route}^r is the regulation component due to (possible) routing of customers to non-basic servers, and functions $Y_{route,i}^r$ representing the contributions into Y_{route}^r due to different flows $i \in I$.

The regulation component $Y_{idle}^r(t)$ is clearly non-negative non-decreasing. It is easy to verify that, each regulation component $Y_{route,i}^r(t)$ is also non-negative and non-decreasing. (And such is, therefore, $Y_{route}^r(t)$.) Moreover, $Y_{route,i}^r(t)$ is a piece-wise constant function, which is constant between type i customer arrivals, and jumps by the value $\alpha_j^* / \mu_{ij} - \nu_i^* \geq 0$ when a type i arrival is routed to server j ; note that the size of the jump is strictly positive if and only if server j is non-basic for type i . Finally, note that $Y^r(t)$ does not increase over some time interval if and only if, during that interval, none of the servers idles and all new arrivals are routed to the corresponding basic servers.

We record the above facts (along with their obvious generalization) in the following lemma for future reference.

Lemma 4 *For each value of the scaling parameter r , consider a pair of time points $0 \leq t_1^r < t_2^r < \infty$, and denote*

$$B_0^r \doteq \frac{Y^r(t_2^r) - Y^r(t_1^r)}{t_2^r - t_1^r} ,$$

$$B_{1,i}^r \doteq \sum_{j \notin J_i} \frac{F_{ij}^r(t_2^r) - F_{ij}^r(t_1^r)}{t_2^r - t_1^r} ,$$

$$B_{2,j}^r \doteq \sum_{i \in I} \frac{G_{ij}^r(t_2^r) - G_{ij}^r(t_1^r)}{t_2^r - t_1^r} .$$

Then, $B_0^r = 0$ if and only if $B_{1,i}^r = 0$ for all i and $B_{2,j}^r = 1$ for all j . Also, $\lim_{r \rightarrow \infty} B_0^r = 0$ if and only if $\lim_{r \rightarrow \infty} B_{1,i}^r = 0$ for all i and $\lim_{r \rightarrow \infty} B_{2,j}^r = 1$ for all j .

Let us consider the process $Z^r = (U^r, {}^u X^r, F^r, {}^{\Sigma} V^r, {}^s N^r, N^r, G^r, H^r, Y^r, Y_{idle}^r, Y_{route}^r)$, where

$$\begin{aligned} U^r &= (U_j^r(t), \quad t \geq 0, \quad j \in J), \\ {}^u X^r &= ({}^u X^r(t), \quad t \geq 0), \\ F^r &= (F_i^r(t), \quad t \geq 0, \quad i \in I), \\ {}^{\Sigma} V^r &= ({}^{\Sigma} V_{ij}^r(l), \quad l \geq 0, \quad i \in I, \quad j \in J), \\ {}^s N^r &= ({}^s N_{ij}^r(l), \quad l \geq 0, \quad i \in I, \quad j \in J), \\ N^r &= (N_{ij}^r(l), \quad l \geq 0, \quad i \in I, \quad j \in J), \\ G^r &= (G_{ij}^r(t), \quad t \geq 0, \quad i \in I, \quad j \in J), \\ H^r &= (H_{ij}^r(t), \quad t \geq 0, \quad i \in I, \quad j \in J), \\ Y^r &= (Y^r(t), \quad t \geq 0), \\ Y_{idle}^r &= (Y_{idle}^r(t), \quad t \geq 0), \\ Y_{route}^r &= (Y_{route}^r(t), \quad t \geq 0). \end{aligned}$$

For each r consider the *fluid scaled* process

$$\Gamma^r Z^r \doteq z^r = (u^r, {}^u x^r, f^r, {}^{\Sigma} v^r, {}^s n^r, n^r, g^r, h^r, y^r, y_{idle}^r, y_{route}^r) ,$$

where the fluid scaling operator Γ^r is applied componentwise, and acts on a scalar function $\Xi = (\Xi(t), \quad t \geq 0)$ as follows:

$$(\Gamma^r \Xi)(t) \doteq \frac{1}{r} \Xi(rt) .$$

Definition 4 A fixed set of functions $z = (u, {}^u x, f, {}^{\Sigma} v, {}^s n, n, g, h, y, y_{idle}, y_{route})$ will be called a *fluid sample path (FSP)* if there exists a sequence \mathcal{R}_f of values of r , and a sequence of sample paths (of the corresponding processes) $\{z^r\}$ such that, as $r \rightarrow \infty$ along the sequence \mathcal{R}_f ,

$$z^r \rightarrow z, \quad u.o.c. ,$$

and, in addition,

$$\|u(0)\| < \infty ,$$

$$(f_i^r(t), t \geq 0) \rightarrow (\lambda_i t, t \geq 0) \quad u.o.c. , \quad i \in I, \quad (56)$$

$$({}^{\Sigma} v_{ij}^r(t), t \geq 0) \rightarrow (\mu_{ij}^{-1} t, t \geq 0) \quad u.o.c. , \quad i \in I, \quad j \in J . \quad (57)$$

Remark. A sequence \mathcal{R}_f existence of which is required in the above definition, may be completely unrelated to the sequence \mathcal{R} we introduced earlier in the definition of the heavy traffic regime.

The following lemma establishes some basic properties of fluid sample paths. We omit the simple proof, which is a direct consequence of the definitions involved.

Lemma 5 *For any fluid sample path z , all its component functions are Lipschitz continuous and, in addition,*

$$\begin{aligned} f_i(t) &= \lambda_i t, \quad t \geq 0, \quad i \in I, \\ {}^\Sigma v_{ij}(t) &= \mu_{ij}^{-1} t, \quad t \geq 0, \quad i \in I, \quad j \in J, \\ {}^s n_{ij}(t) &= (\mu_{ij} \phi_{ij} / \lambda_i) t, \quad t \geq 0, \quad i \in I, \quad j \in J, \\ u_j(t) &= u_j(0) + \sum_i \mu_{ij}^{-1} n_{ij}(\lambda_i t) - \sum_i g_{ij}(t), \quad t \geq 0, \quad j \in J, \\ {}^u x(t) &= \alpha^* \cdot u(t) = {}^u x(0) + y(t), \quad t \geq 0. \end{aligned}$$

Furthermore, both $y(\cdot)$ and ${}^u x(\cdot)$ are non-decreasing (with $y(0) = 0$).

Since all component functions of an FSP are Lipschitz, they are absolutely continuous, and therefore for almost all points $t \in R_+$ (with respect to the Lebesgue measure) the following property holds:

each component function of z has (finite) first derivative at t , and each function $n_{ij}(\cdot)$ has (finite) first derivative at $\lambda_i t$.

We refer to such time points t as *regular*. We adopt a convention that $t = 0$ is *not* a regular point (i.e., in the definition of regular points, we require that proper derivatives exist).

The dynamics of $u(t)$ satisfies the following differential equation and additional conditions at every regular point t :

$$\frac{d}{dt} u(t) = \rho^{in}(t) - \rho^{out}(t), \quad (58)$$

where the components of the J -dimensional vectors $\rho^{in}(t)$ and $\rho^{out}(t)$ are defined as

$$\rho_j^{in}(t) \doteq \sum_i \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) \in \mathcal{K}, \quad (59)$$

$$\rho_j^{out}(t) \doteq \sum_i g'_{ij}(t) \in [0, 1], \quad (60)$$

and for ρ^{out} we have

$$\rho_j^{out}(t) = 1 \quad \text{if } u_j(t) > 0. \quad (61)$$

9.2 Uniform Attraction of Fluid Sample Paths.

For $u \in R_+^J$, denote

$${}^*A(u) \doteq \max_j C'_j(u_j)/\alpha_j^* , \quad {}_*A(u) \doteq \min_j C'_j(u_j)/\alpha_j^* ;$$

$\Phi(u) \doteq 1 - {}_*A(u)/{}^*A(u)$ if $u \neq 0$, and $\Phi(0) \doteq 0$ by convention.

Consider the following functions associated with a fixed FSP. First define

$$J^*(t) = \{j \in J \mid C'_j(u_j(t))/\alpha_j^* = {}^*A(u(t))\}$$

and similarly $J_*(t)$ (with *A replaced by ${}_*A$). Next introduce

$${}^*u_j(t) \doteq \{\zeta \geq 0 \mid C'_j(\zeta)/\alpha_j^* = {}^*A(u(t))\} ,$$

and note that ${}^*u_j(t)$ is well defined since each function $C'_j(\cdot)$ is strictly increasing continuous. Let ${}^*x(t) \doteq \alpha^* \cdot {}^*u(t)$, where ${}^*u(t) = ({}^*u_1(t), \dots, {}^*u_J(t))$, and note that ${}^*x(t) \leq {}^*x(t)$ for all $t \geq 0$.

Finally, note that, at any time t , the following five conditions for $u(t)$ are all equivalent:

$u(t)$ is a fixed point;

${}^*A(u(t)) = {}_*A(u(t))$;

$\Phi(u(t)) = 0$;

${}^*x(t) = {}^*x(t)$;

${}^*u(t) = u(t)$.

The following sequence of lemmas establishes further properties of fluid sample paths, which are less obvious than the basic properties of Lemma 5. The form of the MinDrift rule is used in the proofs in an essential way.

Lemma 6 *Consider a fixed FSP z . Suppose $t > 0$ is a regular point and $u(t) \neq 0$. Then the following properties hold at this t :*

(i) $u_j(t) > 0$ for all $j \in J$.

(ii) We have

$$\sum_{j \in J^*(t)} \alpha_j^* u'_j(t) \leq 0 , \quad \sum_{j \in J_*(t)} \alpha_j^* u'_j(t) \geq 0 . \quad (62)$$

(iii) Moreover, there exists a constant $\epsilon_1 > 0$, which depends on system parameters only, such that if, in addition, $u(t)$ is not a fixed point (i.e. ${}^*A(u(t)) > {}_*A(u(t))$), then

$$\sum_{j \in J^*(t)} \alpha_j^* u'_j(t) \leq -\epsilon_1 , \quad \sum_{j \in J_*(t)} \alpha_j^* u'_j(t) \geq \epsilon_1 . \quad (63)$$

Proof. Let us first prove (iii). Thus, consider regular time point $t > 0$ and suppose that ${}^*A(u(t)) > {}_*A(u(t))$.

The following observation is true:

$$\text{If } i \in I \text{ and } j \in J^*(t) \setminus J_i, \text{ then } n'_{ij}(\lambda_i t) = 0. \quad (64)$$

Indeed, according to the MinDrift-rule and Lemma 3(ii)(c), for all sufficiently large r , the prelimit path z^r is such that in a small interval $[t, t + \epsilon]$, $\epsilon > 0$, new arriving customers of type i cannot be routed to a server $j \in J^*(t) \setminus J_i$. This easily implies that the corresponding FSP component $n'_{ij}(\cdot)$ cannot increase in a small interval to the right of $\lambda_i t$, and therefore $n'_{ij}(\lambda_i t) = 0$ since t is regular. Using similar argument, it is also easy to prove the following property:

$$\text{If } i \in I, J_i \setminus J^*(t) \neq \emptyset, \text{ and } j \in J^*(t), \text{ then } n'_{ij}(\lambda_i t) = 0. \quad (65)$$

Let us denote by $I^*(t)$ the (non-empty) subset of types i such that $J_i \cap J^*(t) \neq \emptyset$. Since graph $\mathcal{G}(\phi) = \mathcal{G}^*$ is connected, there exists at least one $i \in I^*(t)$ such that $J_i \setminus J^*(t) \neq \emptyset$, in which case (by (65)) we have strict inequality:

$$\sum_{j \in J^*(t)} \alpha_j^* \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) = 0 < \sum_{j \in J^*(t)} \alpha_j^* \phi_{ij}. \quad (66)$$

If $i \in I^*(t)$ and $J_i \setminus J^*(t) = \emptyset$, i.e., $J_i \subseteq J^*(t)$, then

$$\sum_{j \in J^*(t)} \alpha_j^* \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) \leq \sum_{j \in J^*(t)} \alpha_j^* \phi_{ij}. \quad (67)$$

Indeed, using (64), the fact that $\alpha_j^* \mu_{ij}^{-1}$ is same across all $j \in J_i$, $\sum_{j \in J} n'_{ij}(\lambda_i t) \leq 1$, and $\sum_{j \in J_i} \mu_{ij} \phi_{ij} / \lambda_i = 1$, we can write:

$$\begin{aligned} \sum_{j \in J^*(t)} \alpha_j^* \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) &= \sum_{j \in J_i} \alpha_j^* \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) \\ &\leq \sum_{j \in J_i} \alpha_j^* \mu_{ij}^{-1} \lambda_i \frac{\mu_{ij} \phi_{ij}}{\lambda_i} = \sum_{j \in J_i} \alpha_j^* \phi_{ij} = \sum_{j \in J^*(t)} \alpha_j^* \phi_{ij}. \end{aligned}$$

As a corollary from (64), we also obtain the following property:

$$\text{If } i \notin I^*(t) \text{ and } j \in J^*(t), \text{ then } n'_{ij}(\lambda_i t) = 0. \quad (68)$$

We will now show that

$$\sum_{j \in J^*(t)} \alpha_j^* u'_j(t) \leq -\epsilon \quad (69)$$

where $\epsilon > 0$ depends only on the subset $J^*(t)$. Indeed,

$$\sum_{j \in J^*(t)} \alpha_j^* u'_j(t) = \sum_{j \in J^*(t)} \alpha_j^* \sum_{i \in I} \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) - \sum_{j \in J^*(t)} \alpha_j^* ,$$

and (using (68), (66) and (67)) we have:

$$\begin{aligned} \sum_{j \in J^*(t)} \alpha_j^* \sum_{i \in I} \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) &= \sum_{i \in I^*(t)} \sum_{j \in J^*(t)} \alpha_j^* \mu_{ij}^{-1} \lambda_i n'_{ij}(\lambda_i t) \\ &< \sum_{i \in I^*(t)} \sum_{j \in J^*(t)} \alpha_j^* \phi_{ij} = \sum_{j \in J^*(t)} \alpha_j^* \sum_{i \in I^*(t)} \phi_{ij} \leq \sum_{j \in J^*(t)} \alpha_j^* . \end{aligned} \quad (70)$$

We have proved (69), with $\epsilon > 0$ depending only on the subset $J^*(t) \subset J$. Since there is only a finite number of subsets of J , we have proved the first inequality in (63), with some fixed $\epsilon_1 > 0$.

The second inequality in (63) is proved analogously. We denote by $I_*(t)$ the (non-empty) subset of types i such that $J_i \cap J_*(t) \neq \emptyset$. Then we use the following property (obtained using the argument analogous to that leading to (64) and (65)):

$$\text{If } i \in I_*(t) \text{ and } j \notin J_i \cap J_*(t), \text{ then } n'_{ij}(\lambda_i t) = 0 .$$

We omit details.

The proof of the non-strict inequalities in property (ii) is a straightforward extension of the proof of (iii). Namely, we need to consider an additional (degenerate) case when $J^*(t) = J_*(t) = J$. In this case, for example to prove the first inequality in (62), we observe that the non-strict inequality (67) always applies, and (70) holds with the strict inequality replaced by a non-strict one.

Finally, (i) is proved by contradiction. Suppose, $u_j(t) = 0$ for some $j \in J$. Obviously, the set of such j is exactly $J_*(t)$. Since $u(t) \neq 0$, $u(t)$ is not a fixed point. Therefore, the second inequality in (63) should hold. However, this is impossible, because we must have $u'_j(t) = 0$ for all $j \in J_*(t)$. Indeed, the condition $u_j(t) = 0$ and the existence of $u'_j(t)$ imply that $u'_j(t) = 0$. (Otherwise $u_j(\cdot)$ would be negative just before or right after time t .) \blacksquare

Lemma 7 Consider a fixed FSP. Suppose a time interval $[t_1, t_2]$, with $0 \leq t_1 < t_2$, is such that

$$\min_{t_1 \leq t \leq t_2} \min_{j \in J} u_j(t) > 0 .$$

Then, over $[t_1, t_2]$, the functions ${}^*A(q(t))$, ${}_*A(q(t))$, ${}^*x(t)$, and ${}^*u_j(t)$ for all $j \in I$, are Lipschitz continuous. Moreover, for almost all $t \in [t_1, t_2]$,

$$\frac{d}{dt} [{}^*A(u(t))] \leq 0, \quad \frac{d}{dt} [{}_*A(u(t))] \geq 0, \quad \frac{d}{dt} [{}^*x(t)] \leq 0, \quad (71)$$

and if, in addition, ${}^*A(u(t)) > {}_*A(u(t))$ (i.e. $u(t)$ is not a fixed point), then

$$\frac{d}{dt}[{}^*x(t)] \leq -\epsilon_1, \quad (72)$$

where $\epsilon_1 > 0$ is defined in Lemma 6.

Proof. First of all, the Lipschitz continuity of each function $C'_j(u_i(t))$ in $[t_1, t_2]$ follows from Lipschitz continuity of $u_j(\cdot)$ and the fact that, for the range of possible values of $u_j(t)$ in $[t_1, t_2]$, $C''_j(\cdot)$ is continuous bounded away from both infinity and 0. (This is the only place where we use the assumption that the functions $C_i(\cdot)$ are twice continuously differentiable.)

This implies that for an arbitrary fixed subset $\hat{J} \subseteq J$, the following functions are also Lipschitz continuous in $[t_1, t_2]$:

$$\max_{i \in \hat{J}} C'_j(u_j(t))/\alpha_j^*, \quad \min_{i \in \hat{J}} C'_j(u_j(t))/\alpha_j^*.$$

In particular, ${}^*A(q(t))$ and ${}_*A(q(t))$ are Lipschitz, which (along with the fact that the second derivatives $C''_j(\cdot)$ are bounded away from 0) implies that all ${}^*u_j(t)$ and ${}^*x(t)$ are Lipschitz.

We see that almost all points $t \in [t_1, t_2]$ are regular (as defined earlier) and, in addition, are such that all the max and min functions in the last display, for all (non-empty) subsets $\hat{J} \subseteq J$, have derivatives. Within the present proof, let us call such points t *strictly regular*.

Consider an arbitrary strictly regular point $t \in [t_1, t_2]$. The proof will be complete once we prove (71) and (72) for this point t .

Since t is strictly regular, the derivatives $d/dt[{}^*A(u(t))]$ and $d/dt[C'_j(u_j(t))/\alpha_j^*]$ for $j \in J^*(t)$, are all equal. (In particular, this implies that ${}^*u'_j(t) = u'_j(t)$ for all $j \in J^*(t)$.)

We cannot have $d/dt[{}^*A(u(t))] > 0$ because this would imply that $u'_j(t) > 0$ for all $j \in J^*(t)$, which would contradict (62). This proves the first (and with it the last) inequality in (71). The second inequality in (71) is proved analogously.

We can now write

$$\frac{d}{dt}[{}^*x(t)] = \sum_{j \in J} \alpha_j^* {}^*u'_j(t) \leq \sum_{j \in J^*(t)} \alpha_j^* {}^*u'_j(t) = \sum_{j \in J^*(t)} \alpha_j^* u'_j(t),$$

where the inequality follows from the fact that ${}^*u'_j(t) \leq 0$ for all $j \in J$ (which is implied by (71)), and the second equality is because ${}^*u'_j(t) = u'_j(t)$ for $j \in J^*(t)$. In the case ${}^*A(q(t)) > {}_*A(q(t))$, by (63), the RHS of the above display is bounded above by $-\epsilon_1$, which proves (72). \blacksquare

Lemma 8 Consider a fixed FSP z . Suppose $u(t_1) \neq 0$ for some $t_1 \geq 0$. Then, $u(t)$ has all strictly positive components (i.e., $u(t) \in R_{++}^J$) for all $t > t_1$. Moreover, in $[t_1, \infty)$, ${}_*A(q(t))$ is non-decreasing, and both ${}^*A(q(t))$ and ${}^*x(t)$ are non-increasing.

Proof. Indeed, we can always find a regular point $\xi > t_1$ arbitrarily close to t_1 so that $u(\xi) \neq 0$. By Lemma 6, $u(\xi) \in R_{++}^J$. Then, using Lemma 7, it follows that ${}_*A(u(t))$ is non-decreasing (and ${}^*A(u(t))$ and ${}^*x(t)$ are non-increasing) starting from time ξ , and therefore $u(t) \in R_{++}^J$ for all $t \geq \xi$. Since ξ can be chosen arbitrarily close to t_1 , the proof is complete. ■

Lemma 9 Consider a fixed FSP z . If $u(0) = 0$ then $u(t) = 0$ for all $t \geq 0$.

Proof. Suppose not. By continuity of ${}^*x(\cdot)$, for an arbitrarily $\epsilon > 0$, there exists time $t_1 > 0$ at which ${}^*x(t)$ reaches level ϵ for the first time. Of course, $u(t_1) \neq 0$. By Lemma 8, ${}^*x(t)$ can not increase starting at time t_1 , and therefore ${}^*x(t) \leq \epsilon$ for all $t \geq 0$. Since $\epsilon > 0$ can be chosen arbitrarily small, ${}^*x(t) = 0$, and therefore $u(t) = 0$ for all $t \geq 0$. ■

The following theorem easily follows from the lemmas presented above in this subsection.

Theorem 5 For any fluid sample path, $\Phi(u(t))$ is a non-increasing function, and the server workload ${}^u x(t)$ is a non-decreasing function. Moreover, there exist fixed constants $T_1 > 0$ and $K \geq 1$ such that, for any FSP, $u(t)$ reaches a fixed point ${}^{\circ}u$ within finite time ${}^u x(0)T_1$ and then stays there, and $\alpha^* \cdot {}^{\circ}u \leq {}^u x(0)K$.

Proof. The fact that $x(t)$ is non-decreasing has already been established earlier.

Suppose $u(0) \neq 0$. By Lemma 8, ${}_*A(u(t))$ is non-decreasing and ${}^*A(u(t))$ is non-increasing in $[0, \infty)$, and therefore $\Phi(u(t))$ is non-increasing. Further, by Lemma 8, $u(t) \in R_{++}^J$ for all $t > 0$. Then, by Lemma 7, for almost all $t > 0$, ${}^*x(t) > {}^u x(t)$ implies

$${}^*x'(t) \leq -\epsilon_1 .$$

Since ${}^*x(0) \leq {}^u x(0)[\sum_j \alpha_j^*]/[\min_j \alpha_j^*]$, ${}^u x(t) \leq {}^*x(t)$, and ${}^u x(t)$ is non-decreasing, we immediately see that $u(t)$ must reach a fixed point within a time proportional to ${}^u x(0)$.

Therefore, the statement of the theorem, with some fixed $T_1 > 0$ and $K \geq 1$, holds for the FSPs with $u(0) \neq 0$. By Lemma 9, it trivially holds for $u(0) = 0$ as well. ■

For future reference, we record the following property of prelimit paths.

Lemma 10 There exists a constant $\epsilon_2 > 0$, such that the following holds. For any prelimit (scaled) path $u^r = (u^r(t), t \geq 0)$, and $0 \leq t_1^r < t_2^r < \infty$, the property

$$u^r(t) \neq 0 \text{ and } \Phi(u^r(t)) \leq \epsilon_2, \quad \forall t \in [t_1^r, t_2^r] ,$$

implies that $y^r(t_2^r) - y^r(t_1^r) = 0$ or, equivalently, that in the (scaled) interval $(t_1^r, t_2^r]$ all new arriving customers are routed to their corresponding basic servers and none of the servers idles.

Proof. First, since $u^r(t) \neq 0$ in $[t_1^r, t_2^r]$, none of the servers idles in this time interval. Second, small value of $\Phi(u^r(t))$ for implies that the vector $(C'_1(u_1^r(t)), \dots, C'_J(u_J^r(t)))$ is “almost proportional” to vector α^* . Thus, if $\Phi(u^r(t))$ is small, it follows from the form of MinDrift(U) rule and Lemma 3(ii)(c) that in $[t_1^r, t_2^r]$ a new arrival of any type $i \in I$ can only be routed to a server $j \in J_i$. We omit the ϵ - δ formalities. \blacksquare

10 Proof of Theorem 1

For each $r \in \mathcal{R}$ consider the following process, obtained by diffusion scaling:

$$\begin{aligned} \tilde{\Gamma}^r(U^r, {}^u X^r, F^r, {}^{\Sigma} V^r, {}^s N^r, N^r, G^r, H^r, Y^r, Y_{idle}^r, Y_{route}^r) \doteq \\ (\tilde{u}^r, {}^u \tilde{x}^r, \tilde{f}^r, {}^{\Sigma} \tilde{v}^r, {}^s \tilde{n}^r, \tilde{n}^r, \tilde{g}^r, \tilde{h}^r, \tilde{y}^r, \tilde{y}_{idle}^r, \tilde{y}_{route}^r), \end{aligned}$$

where the diffusion scaling operator $\tilde{\Gamma}^r$ is defined in (36).

To prove properties stated in Theorem 1, it will suffice to show that for any subsequence $\mathcal{R}_1 \subseteq \mathcal{R}$ there exists another subsequence $\mathcal{R}_2 \subseteq \mathcal{R}_1$ such that these properties hold when $r \rightarrow \infty$ along \mathcal{R}_2 . As in [14], to do this, we will choose subsequence \mathcal{R}_2 and construct all processes (for all $r \in \mathcal{R}_2$) on the same probability space in a way such that the desired properties hold with probability 1 (or are implied by a certain probability 1 properties).

Let us fix an arbitrary subsequence $\mathcal{R}_1 \subseteq \mathcal{R}$ of indices $\{r\}$. According to Skorohod’s representation theorem (see, for example, [7]), for each i , the sequences (on r) of the processes $\{F_i^r\}$ and $\{{}^{\Sigma} V_{ij}^r\}$, $j \in J$, can be constructed on a probability space such that the convergence in (31) holds u.o.c. with probability 1 (w.p.1):

$$\{r^{-1}(A_i^r(r^2 t) - \lambda_i^r \frac{\sum_j \phi_{ij} \alpha_j^*}{\lambda_i} r^2 t), t \geq 0\} \xrightarrow{u.o.c.} \{{}^u \sigma_i B_i(t), t \geq 0\}, \quad (73)$$

where B_i is a standard Brownian motion.

We can and do assume that our underlying probability space $\Omega = \{\omega\}$ is a direct product of the above I probability spaces. (Without loss of generality we assume that this probability space is complete.) On this probability space the convergence (33) holds u.o.c. w.p.1 as well:

$$\{\sum_i r^{-1}(A_i^r(r^2 t) - [\sum_j \alpha_j^*] r^2 t), t \geq 0\} \xrightarrow{u.o.c.} \{at + \sigma B(t), t \geq 0\}, \quad (74)$$

where B is a standard Brownian motion.

Now, from condition (25) and Bramson’s weak law estimates ([4], Proposition 4.3), we know that for any $T_3 > 0$, any $\epsilon > 0$ and any i , for all large r , we have (see the proof of property (5.19) in Proposition 5.1 of [4]):

$$P\{\max_{0 \leq l \leq T_3 r} \sup_{0 \leq \xi \leq 1} |f_i^r(l + \xi) - f_i^r(l) - \lambda_i \xi| \geq \epsilon\} < \epsilon. \quad (75)$$

(The max in (75) and (76), as well as below in (76), (77) and (78), is over *integers* $l \in [0, T_3 r]$.) Also, using Proposition 4.2 of [4], it is easy to show (similarly to the derivation of property (71) in [14]), that for any $T_3 > 0$, any $\epsilon > 0$ and any pair of (i, j) , for all large r , we have

$$P\left\{\max_{0 \leq l \leq T_3 r} \sup_{0 \leq \xi \leq 1} |\Sigma v_{ij}^r(l + \xi) - \Sigma v_{ij}^r(l) - \mu_{ij}^{-1} \xi| \geq \epsilon\right\} < \epsilon. \quad (76)$$

Estimates (75) and (76) enable us to choose a subsequence $\mathcal{R}_2 \subseteq \mathcal{R}_1$, such that, as $r \rightarrow \infty$ along \mathcal{R}_2 , with probability 1, for any $T_3 > 0$ we have

$$\max_{0 \leq l \leq T_3 r} \sup_{0 \leq \xi \leq 1} |f_i^r(l + \xi) - f_i^r(l) - \lambda_i \xi| \rightarrow 0, \quad i \in I, \quad (77)$$

and

$$\max_{0 \leq l \leq T_3 r} \sup_{0 \leq \xi \leq 1} |\Sigma v_{ij}^r(l + \xi) - \Sigma v_{ij}^r(l) - \mu_{ij}^{-1} \xi| \rightarrow 0, \quad i \in I, \quad j \in J. \quad (78)$$

Properties (77) and (78) in turn imply the following property.

With probability 1, for any fixed $T_4 > 0$ and $d > 0$, for any (i, j) , we have:

uniformly on any sequence of pairs (t_1^r, t_2^r) , $r \in \mathcal{R}_2$, such that $0 \leq t_1^r < t_2^r \leq r^2 T_4$, $t_2^r - t_1^r \geq rd$,

$$\lim_{r \rightarrow \infty, r \in \mathcal{R}_2} \frac{\Sigma V_{ij}^r(s N_{ij}^r(F_i^r(r^2 t_2))) - \Sigma V_{ij}^r(s N_{ij}^r(F_i^r(r^2 t_1)))}{\phi_{ij}(t_2^r - t_1^r)} = 1; \quad (79)$$

uniformly on any sequence of pairs (l_1^r, l_2^r) , $r \in \mathcal{R}_2$, such that $0 \leq l_1^r < l_2^r \leq r^2 T_4$, $l_2^r - l_1^r \geq rd$,

$$\lim_{r \rightarrow \infty, r \in \mathcal{R}_2} \frac{\Sigma V_{ij}^r(l_2^r) - \Sigma V_{ij}^r(l_1^r)}{\mu_{ij}^{-1}(l_2^r - l_1^r)} = 1. \quad (80)$$

For each $j \in J$ we have

$$U_j^r(r^2 t) = U_j^r(0) + \sum_i \Sigma V_{ij}^r(N_{ij}^r(F_i^r(r^2 t))) - \sum_i G_{ij}^r(r^2 t),$$

and therefore the expression for the scaled server workload can be written as follows:

$${}^u \tilde{x}^r(t) = \quad (81)$$

$${}^u \tilde{x}^r(0) + r^{-1} \left[\sum_j \alpha_j^* \sum_i \Sigma V_{ij}^r(s N_{ij}^r(F_i^r(r^2 t))) - \sum_j \alpha_j^* r^2 t \right] + \quad (82)$$

$$r^{-1} \sum_j \alpha_j^* (r^2 t - \sum_i G_{ij}^r(r^2 t)) + \quad (83)$$

$$r^{-1} \sum_j \sum_i \alpha_j^* [\Sigma V_{ij}^r (N_{ij}^r (F_i^r (r^2 t))) - \Sigma V_{ij}^r ({}^s N_{ij}^r (F_i^r (r^2 t)))] \quad (84)$$

$$= \tilde{w}^r(t) + \tilde{y}_{idle}^r(t) + \tilde{y}_{route}^r(t) = \tilde{w}^r(t) + \tilde{y}^r(t) , \quad (85)$$

where $\tilde{w}^r(t)$ is the term (82), $\tilde{y}_{idle}^r(t)$ denotes the term (83), $\tilde{y}_{route}^r(t)$ denotes the term (84), and $\tilde{y}^r(t) \doteq \tilde{y}_{idle}^r(t) + \tilde{y}_{route}^r(t)$. We know from (74) that

$$(\tilde{w}^r(t), t \geq 0) \xrightarrow{u.o.c.} (\tilde{w}(t), t \geq 0) ,$$

where

$$\tilde{w}(t) \doteq \tilde{w}(0) + at + \sigma B(t) ,$$

$B(\cdot)$ is the realization of a standard Brownian motion, and the parameters a and σ are those defined in (34). (The realization $\tilde{w}(\cdot)$ is of course continuous.) As seen from (85), the key step in proving Theorem 1, will be the proof of the following convergence:

$$(\tilde{y}^r(t), t \geq 0) \xrightarrow{u.o.c.} (\tilde{y}(t), t \geq 0) . \quad (86)$$

In the rest of this section, we restrict ourselves to a (measurable, probability 1) subset $\Omega_2 \subseteq \Omega$ of elementary outcomes ω , such that all the specified above probability 1 properties hold, when $r \rightarrow \infty$ along \mathcal{R}_2 .

Lemma 11 *Consider a fixed $\omega \in \Omega_2$. As $r \rightarrow \infty$ along \mathcal{R}_2 , the functions \tilde{y}_{route}^r and \tilde{y}_{route} , and then \tilde{y}^r and \tilde{y}^r , are “asymptotically close” in the following sense. For any fixed $T_4 > 0$, and any fixed $\delta_1 > 0$ and $\delta_2 > 0$, for all sufficiently large r , uniformly on $t \in [0, T_4]$,*

$$(1 - \delta_1) \tilde{y}_{route}^r(t) - \delta_2 \leq \tilde{y}_{route}^r(t) \leq (1 + \delta_1) \tilde{y}_{route}^r(t) + \delta_2 , \quad (87)$$

and then

$$(1 - \delta_1) \tilde{y}^r(t) - \delta_2 \leq \tilde{y}^r(t) \leq (1 + \delta_1) \tilde{y}^r(t) + \delta_2 . \quad (88)$$

The proof of Lemma 11 is analogous to the proof of Lemma 9 in [12]. The key observation here (which follows from (79) and (80)) is that, for fixed $T_4 > 0$ and (arbitrarily small) $d > 0$, if $t \in [0, T_4]$ and $|H_{ij}^r(F_i^r(r^2 t))| \geq rd$, then, for all sufficiently large r , the ratio

$$\frac{\Sigma V_{ij}^r (N_{ij}^r (F_i^r (r^2 t))) - \Sigma V_{ij}^r ({}^s N_{ij}^r (F_i^r (r^2 t)))}{\mu_{ij}^{-1} H_{ij}^r (F_i^r (r^2 t))}$$

is close to 1. We do not present the details. We note that (analogously the the situation with Lemma 9 in [12]) Lemma 11 applies to any service discipline satisfying condition (d0), and the uniqueness of ϕ (in the CRP condition) is used in the proof of Lemma 11 in essential way.

It follows from Lemma 11 that to prove (86), it suffices to prove

$$(\tilde{y}^r(t), t \geq 0) \xrightarrow{u.o.c.} (\tilde{y}(t), t \geq 0) , \quad (89)$$

because $\tilde{y}(\cdot)$ is bounded on finite intervals.

Since regulation \tilde{y}^r is a non-decreasing function (for any r), for any fixed $\omega \in \Omega_2$, from any subsequence $\mathcal{R}_3(\omega) \subseteq \mathcal{R}_2$ (which may depend on ω !) it is always possible to find a further subsequence $\mathcal{R}_4(\omega) \subseteq \mathcal{R}_3(\omega)$ such that

$$\tilde{y}^r \Rightarrow \tilde{y} , \quad (90)$$

where \tilde{y} is some non-decreasing RCLL function. (We will prove that this limit \tilde{y} is indeed the regulation of the one-dimensional Brownian motion defined earlier.) In principle, \tilde{y} may take the values $+\infty$. (In other words, $\tilde{y} \in D([0, \infty), \bar{R})$. Recall that notation “ \Rightarrow ” stands for convergence at every point of continuity of the limit function except maybe the point 0.) We note that (90) implies that

$${}^u \tilde{x}^r \Rightarrow \tilde{x} \doteq \tilde{w} + \tilde{y} , \quad (91)$$

and therefore $\tilde{x}(t) < \infty$ if and only if $\tilde{y}(t) < \infty$.

The following lemma (and its proof) is analogous to Lemma 7 in [14] and Lemma 10 in [12]; it contains key observations which are used in the proof of Theorem 1. The key construction of the proof, which involves “slowing down” the diffusion scaled process to consider a family of processes on the “fluid” time scale, and then exploiting the uniform attraction property of fluid sample paths, is essentially same as that in Section 5 of [4]. This construction is central for establishing SSC in the heavy traffic asymptotic regime for multiclass queueing networks (see [4, 19]).

Lemma 12 *Suppose that the service discipline is such that the routing rule is $\text{MinDrift}(U)$ and scheduling rules at the servers are work-conserving. Suppose, $\omega \in \Omega_2$ and a subsequence $\mathcal{R}_4(\omega) \subseteq \mathcal{R}_2$ are fixed such that, along this subsequence, (90) holds. Suppose, a sequence $\{\tilde{t}^r, r \in \mathcal{R}_4(\omega)\}$ is fixed such that*

$$\tilde{t}^r \rightarrow t' \geq 0 ,$$

and

$${}^u \tilde{x}^r(\tilde{t}^r) \rightarrow C > 0 .$$

Let $\delta > 0$ be fixed, and

$$\epsilon \doteq \sup_{\xi_1, \xi_2 \in [t' - 3\delta, t' + 3\delta] \cap R_+} |\tilde{w}(\xi_1) - \tilde{w}(\xi_2)| < C .$$

Then,

- (a) \tilde{y} (and \tilde{x}) is finite in $[0, t' + \delta]$;
- (b) \tilde{y} does not increase in $(t', t' + \delta]$, i.e., $\tilde{y}(t' + \delta) - \tilde{y}(t') = 0$;
- (c) the following bound holds

$$C - \epsilon \leq \tilde{x}(t) \leq CK + \epsilon , \quad \forall t \in [t', t' + \delta] ,$$

with K defined in Theorem 5;

(d) for any $\delta' > 0$,

$$(\tilde{u}^r(t), t \in [t' + \delta', t' + \delta]) \xrightarrow{u.o.c.} (\tilde{u}(t), t \in [t' + \delta', t' + \delta]) ,$$

where $\tilde{u}(t)$ is the (unique) fixed point such that $\alpha^* \cdot \tilde{u}(t) = \tilde{x}(t)$.

If, in addition, $\tilde{t}^r = t'$ for all r , and $\tilde{u}^r(t') \rightarrow {}^o\tilde{u}$, where ${}^o\tilde{u}$ is a fixed point (necessarily, with $\alpha^* \cdot {}^o\tilde{u} = C$), then

(c') $\tilde{x}(t') = C$ and, consequently, $\tilde{u}(t') = {}^o\tilde{u}$;

(d') the following holds:

$$(\tilde{u}^r(t), t \in [t', t' + \delta]) \xrightarrow{u.o.c.} (\tilde{u}(t), t \in [t', t' + \delta]) .$$

Proof essentially repeats that of Lemma 10 in [12]. For completeness, and since some adjustments are required, we present it here.

Let us consider the functions of interest on the fluid time scale. Namely, consider earlier defined functions ${}^u x^r(t) \equiv {}^u \tilde{x}^r(t/r)$, $y^r(t) \equiv \tilde{y}^r(t/r)$, $t \geq 0$, and similarly defined function w^r and other related ones.

Let us choose a fixed $T > 0$ as follows. Let us fix $\epsilon_3 \in (0, C - \epsilon)$, denote

$$C_3 = (C + \epsilon_3)K + \epsilon + \epsilon_3 ,$$

and fix arbitrary

$$T \geq C_3 T_1 ,$$

where K and T_1 are the constants defined in Theorem 5. As seen below in the proof, C_3 will be the upper bound of ${}^u \tilde{x}^r(\cdot)$ in the interval $[\tilde{t}^r, \tilde{t}^r + \delta]$ or, equivalently, the upper bound of ${}^u x^r(\cdot)$ in the interval $[r\tilde{t}^r, r\tilde{t}^r + r\delta]$. Thus, the choice of the constant T is such that an FSP with initial server workload not exceeding C_3 will converge to a fixed point within time T .

For each integer $l \in [0, 2\delta r/T]$, consider

$${}^u \bar{x}^{r,l}(\xi) \doteq {}^u x^r(r\tilde{t}^r + Tl + \xi), \quad \xi \geq 0 ,$$

and similarly defined ${}^u \bar{w}^{r,l}$, ${}^u \bar{y}^{r,l}$, and other related functions.

Let us fix arbitrary $\epsilon_4 \in (0, \epsilon_2/2)$, where ϵ_2 is defined in Lemma 10. Then the following property holds.

Property 1. For all sufficiently large r , the relation (92) below holds for all integer $l \in [0, 2\delta r/T]$, and the relations (93)-(95) hold for all integer $l \in [1, 2\delta r/T]$:

$$C - \epsilon - \epsilon_3 \leq {}^u \bar{x}^{r,l}(\xi) \leq C_3 \quad \forall \xi \in [0, T], \quad (92)$$

$$\Phi({}^u \bar{u}^{r,l}(\xi)) \leq \epsilon_4 \quad \text{for } \xi = 0 \text{ and } \xi = T, \quad (93)$$

$$\Phi(\bar{u}^{r,l}(\xi)) \leq 2\epsilon_4 \quad \forall \xi \in [0, T], \quad (94)$$

$$\bar{y}^{r,l}(T) - \bar{y}^{r,l}(0) = 0. \quad (95)$$

To prove Property 1, we first observe that (92) must hold for $l = 0$ for all large r , because otherwise we would be able to choose a subsequence of indices r along which the sequence of paths $\bar{z}^{r,0}$ converges to an FSP z with ${}^u\bar{x}(0) = C$ and either ${}^u\bar{x}(\xi) > CK$ or ${}^u\bar{x}(\xi) < C$ for some $\xi \in [0, T]$, which contradicts Theorem 5. Moreover, this observation shows that in fact for all large r ,

$$C - \epsilon_3/2 \leq {}^u\bar{x}^{r,1}(0) \leq (C + \epsilon_3/2)K, \quad (96)$$

and, given our choice of the constant T ,

$$\Phi(\bar{u}^{r,1}(0)) \leq \epsilon_4. \quad (97)$$

Next, suppose Property 1 does not hold. Then, we can choose an (infinite) subsequence of r such that (along this subsequence) $l' = l'(r)$ is well defined as the smallest $l \geq 1$ such that one of the conditions (92)-(95) does not hold. (Note that, by this construction and (97), property (93) always holds for $l = l'$ and $\xi = 0$.) We will show that this construction leads to a contradiction.

Indeed, for all large r , both (92) and (93) hold for $l = l'$ and $\xi = 0$. This follows from the combination of the following facts:

property (96);

$|\tilde{w}(\xi_1) - \tilde{w}(\xi_2)| \leq \epsilon$ as long as $\xi_1, \xi_2 \in [t' - 3\delta, t' + 3\delta] \cap R_+$;

$\tilde{w}^r \rightarrow \tilde{w}$ uniformly in $[t' - 3\delta, t' + 3\delta] \cap R_+$;

property (95) for each $1 \leq l \leq l' - 1$;

the functions \tilde{y}^r and \check{y}^r are asymptotically close (in the sense of Lemma 11).

Since (92) and (93), with $l = l'$ and $\xi = 0$, hold for large r , we see that (93) and (94) hold for $l = l'$ and all large r . (Otherwise, we would be able to choose a subsequence of r along which $\bar{z}^{r,l'}$ converges to an FSP z violating Theorem 5.) Similarly, the lower bound in (92) must hold (for large r) for $l = l'$ and $\xi \in [0, T]$. This, in conjunction with (94) and Lemma 10, means that (95) holds for $l = l'$ (for large r). Finally, this and the argument we already used above to prove bound (92) for $l = l'$, $\xi = 0$, shows that in fact (92) holds for $l = l'$ and all $\xi \in [0, T]$ (for large r). We have proved that (92)-(95) hold for $l = l'(r)$ for all large r . This is a contradiction with the construction of the function $l' = l'(r)$ which proves Property 1.

Property 1 (namely (92)) implies that, for all large r ,

$$C - \epsilon - \epsilon_3 \leq {}^u\bar{x}^{r,l}(\xi) \leq C_3, \quad \xi \in [0, T], \quad 0 \leq l \leq 2\delta r/T.$$

Statements (a)-(c) of the Lemma follow from this estimate.

To prove (d), we first notice that (a), (b) and Lemma 11 imply the following uniform convergence for the workload process:

$$({}^u\tilde{x}^r(t), t \in [t' + \delta', t' + \delta]) \xrightarrow{u.o.c.} (\tilde{x}(t), t \in [t' + \delta', t' + \delta]) . \quad (98)$$

Statement (d) then follows from Property 1, the fact that ϵ_4 can be chosen arbitrarily small, and convergence (98).

To prove properties (c') and (d'), we use the same exact construction. It is easy to see that, under the additional assumptions, all conditions (92)-(95) in Property 1 hold for all integer $l \in [0, 2\delta r/T]$ (including 0). Given this, properties (c') and (d') are proved analogously to properties (c) and (d). We omit details. \blacksquare

The rest of the proof of Theorem 1 repeats that of Theorem 1 in [14], and that of Theorem 1 in [12], virtually verbatim. We reproduce it here, with the necessary minor adjustments, for completeness.

10.1 Proof of Theorem 1(i).

To prove this part it suffices to prove the following

Property 2. *As $r \rightarrow \infty$ (along \mathcal{R}_2), for any $\omega \in \Omega_2$ (i.e. with probability 1), we have the following convergence*

$$(\tilde{y}^r(t), t \geq 0) \xrightarrow{u.o.c.} (\tilde{y}(t), t \geq 0) , \quad (99)$$

where \tilde{y} is defined by (39), and

$$(\tilde{u}^r(t), t \geq 0) \xrightarrow{u.o.c.} (\tilde{u}(t), t \geq 0) , \quad (100)$$

where for each t , $\tilde{u}(t)$ is the fixed point such that $\alpha^* \cdot \tilde{u}(t) = \tilde{x}(t)$.

Proof of Property 2. Let us fix $\omega \in \Omega_2$. As explained earlier, for an arbitrary subsequence $\mathcal{R}_3(\omega) \subseteq \mathcal{R}_2$ there exists another subsequence $\mathcal{R}_4(\omega) \subseteq \mathcal{R}_3(\omega)$ such that the convergence (90) holds along this subsequence. Then, the proof of Property 2 will be complete if we can prove the following statements (for the chosen ω , with $r \rightarrow \infty$ along $\mathcal{R}_4(\omega)$). We recall that, at this point, the function \tilde{y} is just *some* limit function - the fact that it is equal to the function defined by (39) is what needs to be proved in order to establish (100).

Step 1. *The limit function \tilde{y} is finite everywhere in $[0, \infty)$.*

Step 2. *The function \tilde{y} is continuous, and $\tilde{y}(0) = 0$.*

Step 3. *If $\tilde{x}(t) > 0$, then t is not a point of increase of \tilde{y} .*

Step 4. *The function \tilde{y} , defined above as a limit, satisfies equation (39).*

Step 5. *Convergence (100) holds.*

In this proof, we will use the convention that $\tilde{y}(0-) = 0$, $\tilde{w}(0-) = \tilde{x}(0-) = \tilde{w}(0)$. So, the case $\tilde{y}(0) > 0$ will be viewed as a discontinuity of \tilde{y} (and \tilde{x}) at 0.

Also, we will use the notation

$$\epsilon(\delta, t) \doteq \sup_{\xi_1, \xi_2 \in [t-\delta, t+\delta] \cap R_+} |\tilde{w}(\xi_1) - \tilde{w}(\xi_2)| .$$

Proof of Step 1. Suppose the statement does not hold. Denote $t^* = \inf\{t \geq 0 \mid \tilde{y}(t) = \infty\}$. The inf is attained because \tilde{y} is RCLL.

We choose a small δ such that $\delta \in (0, t^*)$ if $t^* > 0$, and arbitrary $\delta > 0$ if $t^* = 0$. Let us fix $\epsilon = \epsilon(4\delta, t^*)$. Then we choose a small $\Delta t \in (0, \delta)$ and a large C such that $C > \tilde{x}(t^* - \Delta t) + \epsilon$ if $t^* > 0$, and $C > \tilde{x}(0-) + \epsilon$ if $t^* = 0$. We define

$$\tilde{t}^r = \min\{t \geq (t^* - \Delta t) \vee 0 \mid {}^u\tilde{x}^r(t) \geq C\} ,$$

and choose a further subsequence of $\{r\}$ such that

$$\tilde{t}^r \rightarrow t' \in [t^* - \Delta t, t^*] .$$

(We must have $t' \leq t^*$, because the *limit* function $\tilde{y}(t)$, and therefore $\tilde{x}(t)$, is infinite for all $t \geq t^*$.) It is also easy to see (from (77)) that

$${}^u\tilde{x}^r(t) \rightarrow C .$$

The conditions of Lemma 12 are satisfied, and so \tilde{y} is bounded in $[t', t' + \delta]$ - a contradiction, since $t' + \delta > t^*$. Step 1 has been proved.

Proof of Step 2. Suppose the statement does not hold. The contradiction is obtained very similarly to the way it is done in the proof of Step 1. Let t^* be a discontinuity point (the case $t^* = 0$ is included), i.e. $\tilde{y}(t^*-) < \tilde{y}(t^*)$. Since $\tilde{x} = \tilde{w} + \tilde{y}$ and \tilde{w} is continuous, $\tilde{x}(t^*) - \tilde{x}(t^*-) = \tilde{y}(t^*) - \tilde{y}(t^*-)$. There are two possible cases:

- (a) $\tilde{x}(t^*-) > 0$;
- (b) $\tilde{x}(t^*-) = 0$.

Case (a). In this case we must have $t^* > 0$. (Indeed, by the definition of \tilde{w} and our conventions, $\tilde{x}(0-) = \tilde{w}(0) = \lim_r {}^u\tilde{x}^r(0)$. If $\tilde{w}(0) > 0$, then, by Lemma 12(c'), $\tilde{x}(0) = \lim_r {}^u\tilde{x}^r(0)$, which means that \tilde{x} , and therefore \tilde{y} , has no jump at 0. If $\tilde{w}(0) = 0$ then $\tilde{x}(0-) = 0$.) We can always fix a small $\delta > 0$, small $\Delta t \in (0, \delta)$, such that $t' = t^* - \Delta t$ is a point of continuity of \tilde{y} (and \tilde{x}) and $\epsilon = \epsilon(4\delta, t^*) < \tilde{x}(t') = C$. We have convergence ${}^u\tilde{x}^r(t') \rightarrow C$ (since \tilde{x} is continuous at t'), and by Lemma 12, \tilde{y} can not increase in the interval $(t', t' + \delta]$ which contains t_* . So, \tilde{x} can not have a jump at t_* .

Case (b). In this case, let us fix a small $C > 0$ and then a sufficiently small $\delta > 0$ so that

$$C_1 = KC + \epsilon < \tilde{x}(t^*) ,$$

where $\epsilon = \epsilon(4\delta, t^*)$ and $K \geq 1$ is defined in Theorem 5 (and used in Lemma 12). Then, if $t^* > 0$ we fix a small Δt such that

$$\limsup_{r \rightarrow \infty} \sup_{[t^* - \Delta t, t^*]} {}^u \tilde{x}^r(\xi) < C .$$

If $t^* = 0$, we fix an arbitrary $\Delta t > 0$.

We define

$$\tilde{t}^r = \min\{t \geq (t^* - \Delta t) \vee 0 \mid {}^u \tilde{x}^r(t) \geq C\} ,$$

and choose a further subsequence of $\{r\}$ such that

$$\tilde{t}^r \rightarrow t' \in [(t^* - \Delta t) \vee 0, t^*] .$$

The conditions of Lemma 12 are satisfied, and so $\tilde{x}(t) < C_1$ for all $t \in [t', t' + \delta]$, which contradicts the assumption of case (b), since t^* belongs to the latter interval. Step 2 has been proved.

Proof of Step 3. Let $t^* \geq 0$ be such that $\tilde{x}(t^*) > 0$. If $t^* = 0$, then the fact that \tilde{y} does not increase in a small interval $[0, \delta]$ follows from Lemma 12(b'). If $t^* > 0$, then precisely the same construction as in the proof of Step 2(a) shows that \tilde{y} does not increase in a small interval $[t', t' + \delta]$ containing t^* in its interior. Step 3 has been proved.

Proof of Step 4 follows from the statements of Steps 2 and 3, and Proposition 1 (in the Appendix).

Proof of Step 5. It suffices to show that

For any $t^ \geq 0$ and any $\epsilon > 0$, there exists $\delta > 0$ such that*

$$\limsup_{r \rightarrow \infty} \sup_{\xi \in [t^* - \delta, t^* + \delta] \cap R_+} \|\tilde{u}^r(\xi) - \tilde{u}(\xi)\| < \epsilon . \quad (101)$$

(The u.o.c. convergence will then follow from the Heine-Borel lemma.)

If $\tilde{x}(t^*) = 0$, then (101) must hold because both functions \tilde{u} and \tilde{u}^r (for large r) are bounded by an arbitrarily small constant in a sufficiently small neighborhood of t^* . If $\tilde{x}(t^*) > 0$ and $t^* = 0$, then (101) follows from Lemma 12(d'). If $\tilde{x}(t^*) > 0$ and $t^* > 0$, then to obtain (101) we can repeat the construction of the proof of Step 2(a) and then apply Lemma 12(d). Step 5 has been proved.

Thus, the proof of Property 2, and with it the proof of statement (i) of the theorem, is complete. ■

10.2 Proof of Theorem 1(ii).

We use the same construction of the probability space Ω , the subsequence \mathcal{R}_2 , and the probability 1 subset Ω_2 , as specified above. Consider an arbitrary discipline G . Sample

paths for both the $Gc\mu$ and G disciplines are constructed on this common probability space. For $\omega \in \Omega_2$, consider paths of ${}^u\tilde{x}_G^r$, \tilde{y}_G^r , and \tilde{w}_G^r , corresponding to the discipline G . Since \tilde{w}_G^r is invariant with respect to the discipline, $\tilde{w}_G^r = \tilde{w}^r$, and therefore $\tilde{w}_G^r \rightarrow \tilde{w}_G = \tilde{w}$ u.o.c.

We claim that, along the subsequence \mathcal{R}_2 , for any $t \geq 0$,

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [{}^u\tilde{x}_G^r(\xi) - \tilde{x}(\xi)] \geq 0, \quad (102)$$

and therefore (40) holds. To prove this we first recall that *Lemma 11 holds for any discipline G satisfying condition (d0)*.

For any subsequence $\mathcal{R}_3(\omega) \subseteq \mathcal{R}_2(\omega)$, we can choose a further subsequence $\mathcal{R}_4(\omega) \subseteq \mathcal{R}_3(\omega)$ such that $\tilde{y}_G^r \Rightarrow \tilde{y}_G$, where \tilde{y}_G is some non-decreasing non-negative RCLL function. (The case that $\tilde{y}_G(t)$ takes value $+\infty$ starting from some finite time t_* is possible.) Therefore, for any $t \geq 0$ where $\tilde{y}_G(\cdot)$ is continuous, as $r \rightarrow \infty$ along $\mathcal{R}_4(\omega)$,

$$\lim {}^u\tilde{x}_G^r(t) = \tilde{w}(t) + \tilde{y}_G(t).$$

Since ${}^u\tilde{x}_G^r(t)$ is non-negative, we see that $\tilde{w} + \tilde{y}_G$ is non-negative at every point of continuity of \tilde{y}_G , and therefore it is non-negative for all $t \geq 0$ (by right-continuity). Then, by Proposition 1(ii) (in the Appendix), $\tilde{y}_G(t) \geq \tilde{y}(t)$ for all $t \geq 0$. Since \tilde{y} is continuous and non-decreasing, and \tilde{y}_G and all \tilde{y}_G^r are non-decreasing, for any $t \geq 0$ we obtain the uniform bound

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [\tilde{y}_G^r(\xi) - \tilde{y}(\xi)] \geq 0.$$

By Lemma 11, we have analogous bound for \tilde{y}_G^r as well:

$$\liminf_{r \rightarrow \infty} \inf_{\xi \in [0, t]} [\tilde{y}_G^r(\xi) - \tilde{y}(\xi)] \geq 0,$$

which proves (102), with $r \rightarrow \infty$ along subsequence $\mathcal{R}_4(\omega)$, and therefore along \mathcal{R}_2 as well (since the subsequence $\mathcal{R}_3(\omega)$ can be arbitrary). The proof of (102) (and therefore (40)) is complete.

Since the function $\sum_j C_j(u_j)$ is continuous in the vector u , and the fixed point $\tilde{u}(t)$ in (41) minimizes the value of $\sum_j C_j(u_j)$ over vectors u with server workload ${}^u\tilde{x}(t)$, property (41) also holds. Finally, the equality in (42) follows from the fact that $\tilde{u}^r \rightarrow \tilde{u}$ u.o.c., and the inequality follows from (41) and Fatou's lemma.

The proof of Theorem 1 is now complete. ■

11 Proof of Theorem 2

The proof of Theorem 2 is a relatively straightforward extension of that of Theorem 1. The extension is based on the fact that, given the assumptions of Theorem 2(ii), the processes ${}^q\tilde{u}^r$

and \tilde{u}^r are in fact “asymptotically close” (see (103) below). As a result, the behavior of the system (in the diffusion limit) under $\text{MinDrift}(Q)$ is the “same” as that under $\text{MinDrift}(U)$. In this section we provide a detailed sketch of such a proof extension. We believe the details can be easily filled in by a reader.

Construction of the probability space and subsequences \mathcal{R}_1 and \mathcal{R}_2 . For the proof of Theorem 2 we assume that, for each r , the service times of the “initial customers” of type i at server j , whose service has not yet started at initial time 0, are given by an i.i.d. sequence $\bar{v}_{ij}^r(n)$, $n = 1, 2, \dots$. Thus, the sequence $\{\bar{v}_{ij}^r(n)\}$ is separate from the sequence $\{v_{ij}^r(n)\}$, defining service times of customers arriving after time 0, but of course $\bar{v}_{ij}^r(1)$ has the same distribution as $v_{ij}^r(1)$. We denote by

$$\Sigma \bar{V}_{ij}^r(n) \doteq \sum_{m=1}^n \bar{v}_{ij}^r(m), \quad n = 0, 1, 2, \dots,$$

the total amount of unfinished work “contained” in the the first n (in the order of them being taken for service) initial type i customers at the server j . As with other functions, we extend the domain of $\Sigma \bar{V}_{ij}^r(\cdot)$ to all real non-negative $t \geq 0$, and denote its fluid-scaled version by $\Sigma \bar{v}_{ij}^r = \Gamma^r \Sigma \bar{V}_{ij}^r$.

The underlying probability space is the same as in the proof of Theorem 1, except it is augmented by taking a direct product with the space on which the sequences (on r) of the processes $\{\Sigma \bar{V}_{ij}^r\}$ are defined. The subsequence \mathcal{R}_1 is defined exactly same way. The property analogous to (78) holds for the processes $\Sigma \bar{v}_{ij}^r$, as well as Σv_{ij}^r . Then, the subsequence \mathcal{R}_2 can be chosen in a way such that, additionally, the properties analogous to (78) and (80) hold for the processes $\Sigma \bar{v}_{ij}^r$ and $\Sigma \bar{V}_{ij}^r$, respectively.

“Asymptotic closeness” of ${}^q \tilde{u}^r$ and \tilde{u}^r . Using (78) and (80), and their analogs for $\Sigma \bar{v}_{ij}^r$ and $\Sigma \bar{V}_{ij}^r$, it is easy to demonstrate the following property, which holds for any service discipline within the class specified in Theorem 2(ii):

As $r \rightarrow \infty$ along \mathcal{R}_2 , with probability 1, for any $T_3 > 0$ and any $\epsilon > 0$, we have

$$\sup_{0 \leq t \leq T_3} \frac{\|{}^q \tilde{u}^r(t) - \tilde{u}^r(t)\|}{\max(\tilde{u}^r(t), \epsilon)} \rightarrow 0. \quad (103)$$

This property is the key in showing that, in the heavy traffic limit, $\text{MinDrift}(Q)$ induces the same system behavior as the $\text{MinDrift}(U)$.

Definition and properties of the FSPs under $\text{MinDrift}(Q)$. The process Z^r is augmented by the following components:

$$Q^r = (Q_{ij}^r(t), \quad t \geq 0, \quad i \in I, \quad j \in J),$$

$$\Sigma \bar{V}^r = (\Sigma \bar{V}_{ij}^r(l), \quad l \geq 0, \quad i \in I, \quad j \in J),$$

$${}^q U^r = ({}^q U_j^r(t), \quad t \geq 0, \quad j \in J),$$

$${}^q X^r = ({}^q X^r(t), \quad t \geq 0).$$

The fluid-scaled process z^r and an FSP z are augmented by the corresponding components $q^r, \Sigma \bar{v}^r, {}^q u^r, {}^q x^r$, and $q, \Sigma \bar{v}, {}^q u, {}^q x$, respectively. The definition of the FSP is same, except it includes the additional conditions:

$${}^q u(0) = u(0)$$

and an analog of (57) for ${}^q v_{ij}^r$. This augmented definition of an FSP, easily yields the following additional FSP property (which can be added into Lemma 5):

$${}^q u(t) = u(t) \quad \forall t \geq 0, \quad (104)$$

which of course also implies ${}^q x(t) = {}^u x(t)$, $t \geq 0$. Using (104), it can be easily shown that *all* the FSP properties established for MinDrift(U), hold for MinDrift(Q) as well.

Proof of Theorem 2. Given property (103), and the fact that FSPs under MinDrift(Q) satisfy all the properties of FSPs under MinDrift(U) (plus (104)), the rest of the proof is the same as that of Theorem 1.

12 Proof of Theorem 4

Before proceeding with the proof, note that, in addition to (51), we have another pathwise relation (see (21) and (22)):

$${}^q \tilde{x}^r(t) \leq C_0 \tilde{x}^r(t), \quad t \geq 0. \quad (105)$$

Proof. The argument leading to Theorem 3 shows that, given the conditions of Theorem 4, either of the convergences (53) or (54) implies (52). So, it will suffice to prove that (52) implies (55).

Assume (52). Suppose, for some $t_1 \geq 0$ property (55), with $t = t_1$, does not hold. This implies that, in addition to the pathwise inequality $\tilde{x}^r(t_1) \leq {}^q \tilde{x}^r(t_1)$ (see (51)), we have, for some fixed constant $c > 0$,

$$\liminf_{r \rightarrow \infty} P\{{}^q \tilde{x}^r(t_1) - \tilde{x}^r(t_1) > c\} > c. \quad (106)$$

Consider a subsequence of indices r along which the distributions of both $\tilde{x}^r(t_1)$ and ${}^q \tilde{x}^r(t_1)$ (weakly) converge to some distributions, which we denote η and ${}^q \eta$, respectively. Distribution η is necessarily equal to the distribution of $\tilde{x}(t_1)$, and ${}^q \eta$ (stochastically) dominates η , and is *not equal to* η , which follows from (106).

Consider the processes \tilde{x}^r and ${}^q \tilde{x}^r$ restarted at time t_1 . Let us fix arbitrary t_2 , $t_1 < t_2 < \infty$. Then, we have

$$\liminf_r P\{\inf_{[t_1, t_2]} \tilde{x}^r(t) < \epsilon\} \geq p_1 \quad \forall \epsilon > 0, \quad (107)$$

where

$$p_1 = P\{\tilde{x}(t) \text{ hits 0 within } [t_1, t_2]\}.$$

Since pathwise inequality (105) holds, we see that (107) holds for the process ${}^q\tilde{x}^r$ as well:

$$\liminf_r P\left\{\inf_{[t_1, t_2]} {}^q\tilde{x}^r(t) < \epsilon\right\} \geq p_1 \quad \forall \epsilon > 0. \quad (108)$$

Consider now an RBM ${}^q\tilde{x}$ with the drift a and diffusion coefficient σ (same as for the RBM \tilde{x}), defined within interval $[t_1, t_2]$, with the initial distribution ${}^q\eta$ at time t_1 . Since ${}^q\eta$ strictly dominates η ,

$$P\{{}^q\tilde{x}(t) \text{ hits 0 within } [t_1, t_2]\} = p_2 < p_1. \quad (109)$$

Using Theorem 2(ii), it is easy to see that the RBM ${}^q\tilde{x}$ is an asymptotic (stochastic) lower bound of the sequence of processes ${}^q\tilde{x}^r$ (in the sense specified in Theorem 2(ii)). From this fact we see that for any $\delta > 0$ we can choose a sufficiently small $\epsilon > 0$, so that

$$\liminf_r P\left\{\inf_{[t_1, t_2]} {}^q\tilde{x}^r(t) > \epsilon\right\} \geq (1 - p_2) - \delta. \quad (110)$$

If we fix $\delta \in (0, p_1 - p_2)$ and a corresponding $\epsilon > 0$ as above, we obtain from the estimates (110) and (108):

$$\liminf_r \left[P\left\{\inf_{[t_1, t_2]} {}^q\tilde{x}^r(t) > \epsilon\right\} + P\left\{\inf_{[t_1, t_2]} {}^q\tilde{x}^r(t) < \epsilon\right\} \right] \geq p_1 + (1 - p_2) - \delta > 1,$$

a contradiction which completes the proof. ■

13 Stability Vs. Heavy Traffic Workload Minimization

Consider a special case of the MinDrift routing rule, with cost functions $C_j(\zeta) = (1/2)\zeta^2$. (See Section 4.4.) The corresponding MinDrift(Q) rule is: *route an arriving type i customer to a server j such that*

$$j \in \arg \min_{j \in J} {}^qU_j(t)/\mu_{ij}. \quad (111)$$

Our heavy traffic results apply to this rule. (The form of the rule does not change with r , as explained in Section 8).

Using the approach of [15, 6, 2, 1, 14], it is not hard to show that, under this routing rule (plus arbitrary scheduling rules satisfying (d1)-(d2)), both the queue length process $((Q_{ij}(t), i \in I, j \in J), t \geq 0)$ and the unfinished work process $((U_j(t), j \in J), t \geq 0)$ are *stable*, as long as the vector of mean rates λ is within the system stability region \mathcal{M}^0 , defined in Section 5. In fact, as explained in [14], the analysis of the FSPs (Section 9 in

this paper), required to establish the heavy traffic results, is essentially a “superset” of the analysis needed to prove stability. (We do not provide details of the stability proof, as it is not the focus of this paper.)

Thus, the above rule is able to both keep queues stable (as long as $\lambda \in \mathcal{M}^0$) and minimize system workload in the heavy traffic limit. The $Gc\mu$ scheduling rule for the IQ-system possesses the same property (see [12]) and so does the MaxWeight scheduling rule for a different, but closely related, “generalized switch” model (see [14]). All these results may suggest the intuition that a dynamic service discipline which keeps queues stable (as long as $\lambda \in \mathcal{M}^0$), “typically,” will also minimize system workload in heavy traffic. Such a “conjecture” cannot be formally correct, because it is not hard to devise some *contrived* disciplines, for which it does not hold. We note, however, that this conjecture does not hold even for *very natural* service disciplines, as the following example demonstrates.

Consider a service discipline for our OQ-system, which strives to minimize the drift of the cost (Lyapunov) function

$$\sum_{ij} \frac{1}{2} Q_{ij}^2(t).$$

Then, the discipline has the following form. (It is close to the class of network scheduling disciplines introduced in [15].)

Routing rule (“Join the shortest queue of your type”): route an arriving type i customer to a server j such that

$$j \in \arg \min_{j \in J: \mu_{ij} > 0} Q_{ij}(t). \quad (112)$$

Scheduling rule (“ $Gc\mu$ within each server”): server j picks a customer of type i such that

$$i \in \arg \max_{i \in I} Q_{ij}(t) \mu_{ij}. \quad (113)$$

This discipline ensures stability of the queues, when $\lambda \in \mathcal{M}^0$. (Again, the approach and techniques of [15, 6, 2, 1, 14] can be applied.)

However, it is not hard to see that, under this discipline and under the conditions of Theorem 4, condition (55) cannot possibly hold, as long as $\mu_{ij} > 0$ for at least one non-basic activity (ij) . We do not provide a formal proof here. The key part of a proof is showing the following very intuitive fact, which is implied by the nature of the routing rule: *FSPs under this discipline are such that, if the initial workload is non-zero, then after some finite time all non-basic queue lengths are bounded away from 0*. We also exploit the fact that, since the limiting workload process is lower bounded by an RBM, at any time $t > 0$, the limiting workload is non-zero with non-zero probability. Thus, by Theorem 4, *none of the workload minimization properties (52), (53), (54), can hold*.

14 Appendix: The One-Dimensional Skorohod Problem

The following proposition describes standard properties of solutions to the one-dimensional Skorohod problem. (See for example [5] for the proof. The proof is also contained in the proof of Theorem 5.1 of [18]).

Proposition 1 *Let $w = (w(t), t \geq 0)$ be a continuous function in $D([0, \infty), R)$ such that $w(0) \geq 0$. Then the following holds.*

- (i) *There exists a unique pair (x, y) of functions in $D([0, \infty), \bar{R})$, such that*
 - (a) $x(t) = w(t) + y(t) \geq 0$, $t \geq 0$,
 - (b) y is non-decreasing and non-negative,
 - (c) $y(0) = 0$,
 - (d) *for any $t \geq 0$, if $x(t) > 0$ then t is not a point of increase of y , i.e., there exists $\delta > 0$ such that $y(\xi)$ is constant in $[t - \delta, t + \delta] \cap R_+$.*

This unique pair is (x°, y°) , where

$$y^\circ(t) \doteq -[0 \wedge \inf_{0 \leq u \leq t} w(u)], \quad x^\circ(t) = w(t) + y^\circ(t), \quad t \geq 0.$$

- (ii) *For any pair (x, y) of functions in $D([0, \infty), \bar{R})$ satisfying (a) and (b), we have*

$$y(t) \geq y^\circ(t), \quad x(t) \geq x^\circ(t), \quad t \geq 0.$$

Acknowledgement. I'm very grateful to Avi Mandelbaum for bringing output-queued flexible server systems to my attention and encouraging this work.

References

- [1] M. Andrews, K. Kumaran, K. Ramanan, A. L. Stolyar, R. Vijayakumar, P. Whiting. Scheduling in a Queueing System with Asynchronously Varying Service Rates. *Probability in the Engineering and Informational Sciences*, 2004, Vol. 18, pp. 191-217.
- [2] M. Armony, N. Bambos. Queueing Networks with Interacting Service Resources. *Proceedings of the 37th Annual Allerton Conference*, 1999, Urbana, pp. 42-52.
- [3] S.L. Bell, R.J. Williams. Dynamic Scheduling of a System with Two Parallel Servers in Heavy Traffic with Complete Resource Pooling: Asymptotic Optimality of a Continuous Review Threshold Policy. *Annals of Applied Probability*, Vol. 11, (2001), pp. 608-649.
- [4] M. Bramson. State Space Collapse with Applications to Heavy Traffic Limits for Multi-class Queueing Networks. *Queueing Systems*, Vol. 30, (1998), pp. 89-148.

- [5] H. Chen, A. Mandelbaum. Leontief Systems, RBV's and RBM's. In *Applied Stochastic Analysis*, M.H.A.Davis and R.J.Elliott (eds.), Gordon and Breach Science Publishers, New York, 1991, pp. 1-43.
- [6] J.G.Dai, B.Prabhakar. The Throughput of Data Switches with and without Speedup. *Proceedings of the INFOCOM'2000*, pp. 556-564.
- [7] S. N. Ethier and T. G. Kurtz. *Markov Process: Characterization and Convergence*. John Wiley and Sons, New York, 1986.
- [8] J. M. Harrison. Heavy Traffic Analysis of a System with Parallel Servers: Asymptotic Optimality of Discrete Review Policies. *Annals of Applied Probability*, Vol. 8, (1998), pp. 822-848.
- [9] J.M.Harrison, M.J.Lopez. Heavy Traffic Resource Pooling in Parallel-Server Systems. *Queueing Systems*, Vol. 33, (1999), pp. 339-368.
- [10] F. P. Kelly and C. N. Laws. Dynamic Routing in Open Queueing Networks: Brownian Models, Cut Constraints and Resource Pooling. *Queueing Systems*, Vol. 13, (1993), pp. 47-86.
- [11] C. N. Laws. Resource Pooling in Queueing Networks with Dynamic Routing. *Adv. Appl. Prob.*, Vol. 24, (1992), pp. 699-726.
- [12] A. Mandelbaum, A. L. Stolyar. Scheduling Flexible Servers with Convex Delay Costs: Heavy-Traffic Optimality of the Generalized $c\mu$ -rule. *Operations Research*, (2004), to appear.
- [13] N.McKeown, V.Anantharam, and J.Walrand. Achieving 100% Throughput in an Input-Queued Switch. *Proceedings of the INFOCOM'96*, (1996), pp. 296-302.
- [14] A. L. Stolyar. MaxWeight Scheduling in a Generalized Switch: State Space Collapse and Workload Minimization in Heavy Traffic. *Annals of Applied Probability*, 2004, Vol.14, No.1, pp.1-53.
- [15] L. Tassiulas, A. Ephremides. Stability Properties of Constrained Queueing Systems and Scheduling Policies for Maximum Throughput in Multihop Radio Network. *IEEE Transactions on Automatic Control*, Vol. 37, (1992), pp. 1936-1948.
- [16] Y. Teh and A. Ward. Critical Thresholds for Dynamic Routing in Queueing Networks. *Queueing Systems*, Vol. 42, (2002), pp. 297-316.
- [17] J.A.Van Mieghem. Dynamic Scheduling with Convex Delay Costs: the Generalized $c\mu$ Rule. *Annals of Applied Probability*, Vol. 5, (1995), pp. 809-833.
- [18] R.J.Williams. An Invariance Principle for Semimartingale Reflecting Brownian Motions in an Orthant. *Queueing Systems*, Vol. 30, (1998), pp. 5-25.

- [19] R.J.Williams. Diffusion Approximations for Open Multiclass Queueing Networks: Sufficient Conditions Involving State Space Collapse. *Queueing Systems*, Vol. 30, (1998), pp. 27-88.
- [20] R.J.Williams. On Dynamic Scheduling of a Parallel Server System with Complete Resource Pooling. *Fields Institute Communications*, Vol. 28, (2000), pp. 49-71.